

Cervical abnormalities are more common among Indigenous than other Australian women

A retrospective record-linkage study, 2000-2011

Whop, Lisa; Baade, Peter; Garvey, Gail; Cunningham, Joan; Brotherton, Julia; Lokuge, Kamalini; Valery, Patricia ; O'Connell, Dianne L.; Canfell, Karen; Diaz, Abbey; Roder, David; Gertig, DM; Moore, Suzanne; Condon, John

Published in:
PLoS One

DOI:
[10.1371/journal.pone.0150473](https://doi.org/10.1371/journal.pone.0150473)

Published: 01/01/2016

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Whop, L., Baade, P., Garvey, G., Cunningham, J., Brotherton, J., Lokuge, K., Valery, P., O'Connell, D. L., Canfell, K., Diaz, A., Roder, D., Gertig, DM., Moore, S., & Condon, J. (2016). Cervical abnormalities are more common among Indigenous than other Australian women: A retrospective record-linkage study, 2000-2011. *PLoS One*, 11(4), 1-12. Article e0150473. <https://doi.org/10.1371/journal.pone.0150473>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

Cervical Abnormalities Are More Common among Indigenous than Other Australian Women: A Retrospective Record-Linkage Study, 2000–2011

Lisa J. Whop^{1*}, Peter Baade², Gail Garvey¹, Joan Cunningham¹, Julia M. L. Brotherton^{3,4}, Kamalini Lokuge⁵, Patricia C. Valery^{1,6}, Dianne L. O'Connell^{7,8,9}, Karen Canfell^{7,8}, Abbey Diaz¹, David Roder¹⁰, Dorota M. Gertig⁴, Suzanne P. Moore¹, John R. Condon¹

1 Epidemiology and Health Systems Division, Menzies School of Health Research, Charles Darwin University, Darwin, Brisbane, Australia, **2** Cancer Council Queensland, Brisbane, Australia, **3** Victorian Cytology Service Inc, Melbourne, Victoria, Australia, **4** School of Population and Global Health, University of Melbourne, Melbourne, Victoria, Australia, **5** National Centre of Epidemiology and Public Health, Australian National University, Canberra, Australia, **6** QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia, **7** Cancer Research Division, Cancer Council New South Wales, Sydney, New South Wales, Australia, **8** School of Public Health, University of Sydney, Sydney, New South Wales, Australia, **9** School of Medicine and Public Health, University of Newcastle, Newcastle, New South Wales, Australia, **10** Centre for Population Health Research, University of South Australia, Adelaide, South Australia, Australia

* lisa.whop@menzies.edu.au



OPEN ACCESS

Citation: Whop LJ, Baade P, Garvey G, Cunningham J, Brotherton JML, Lokuge K, et al. (2016) Cervical Abnormalities Are More Common among Indigenous than Other Australian Women: A Retrospective Record-Linkage Study, 2000–2011. PLoS ONE 11(4): e0150473. doi:10.1371/journal.pone.0150473

Editor: Marcia Edilaine Lopes Consolaro, State University of Maringá/Universidade Estadual de Maringá, BRAZIL

Received: December 15, 2015

Accepted: February 15, 2016

Published: April 11, 2016

Copyright: © 2016 Whop et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are available in the paper.

Funding: LJ Whop is supported by a Sidney Myer Health Scholarship, Menzies Enhanced Living Top-up and a student scholarship funded by the Lowitja Institute. J Cunningham was supported by a National Health & Medical Research Council (NHMRC) Research Fellowship [#1058244]. PC Valery was supported by a NHMRC Career Development Fellowship [#1083090]. A Diaz is supported by a National Health and Medical Research Council

Abstract

Indigenous Australian women have much higher incidence of cervical cancer compared to non-Indigenous women. Despite an organised cervical screening program introduced 25 years ago, a paucity of Indigenous-identified data in Pap Smear Registers remains. Prevalence of cervical abnormalities detected among the screened Indigenous population has not previously been reported. We conducted a retrospective cohort study of population-based linked health records for 1,334,795 female Queensland residents aged 20–69 years who had one or more Pap smears during 2000–2011; from linked hospital records 23,483 were identified as Indigenous. Prevalence was calculated separately for Indigenous and non-Indigenous women, for cytology-detected low-grade (cLGA) and high-grade abnormalities (cHGA), and histologically confirmed high-grade abnormalities (hHGA). Odds ratios (OR) were estimated from logistic regression analysis. In 2010–2011 the prevalence of hHGA among Indigenous women (16.6 per 1000 women screened, 95% confidence interval [CI] 14.6–18.9) was twice that of non-Indigenous women (7.5 per 1000 women screened, CI 7.3–7.7). Adjusted for age, area-level disadvantage and place of residence, Indigenous women had higher prevalence of cLGA (OR 1.4, CI 1.3–1.4), cHGA (OR 2.2, CI 2.1–2.3) and hHGA (OR 2.0, CI 1.9–2.1). Our findings show that Indigenous women recorded on the Pap Smear Register have much higher prevalence for cLGA, cHGA and hHGA compared to non-Indigenous women. The renewed cervical screening program, to be implemented in 2017, offers opportunities to reduce the burden

(NHMRC) Training Scholarship for Indigenous Australian Health Research (No. 1055587). The National Indigenous Cervical Screening Project is funded by a NHMRC Project Grant [#104559]. This study was undertaken under the auspices of the Centre of Research Excellence in Discovering Indigenous Strategies to improve Cancer Outcomes Via Engagement, Research Translation and Training (DISCOVER-TT CRE, funded by the NHMRC [#1041111]), and the Strategic Research Partnership to improve Cancer control for Indigenous Australians (STREP Ca-CInDA, funded through Cancer Council NSW [SRP 13-01] with supplementary funding from Cancer Council WA). We also acknowledge the ongoing support of the Lowitja Institute, Australia's National Institute for Aboriginal and Torres Strait Islander Health Research. The views expressed in this publication are those of the authors and do not necessarily reflect the views of the funding agencies. Funding bodies have no role in the study design, data collection, access, analysis or interpretation or decision to publish.

Competing Interests: KC, DG and JMLB are investigators on a trial of cytology and primary HPV screening in Australia ('Compass'), which is conducted and funded by the Victorian Cytology Service (VCS), a government-funded health promotion charity and by KC's research group (formerly at UNSW Australia, now at Cancer Council NSW). The VCS has received equipment and a funding contribution for the Compass pilot study and for the main trial from Roche Molecular Systems and Ventana Inc USA. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

of abnormalities and invasive cancer among Indigenous women and address long-standing data deficiencies.

Introduction

The Australian National Cervical Screening Program (NCSP), introduced in 1991, recommends routine two-year screening by Papanicolaou (Pap) smears for women aged 20 to 69 years who have ever been sexually active, regardless of human papillomavirus (HPV) vaccination status, ethnicity, sexual orientation or religion [1]. The major aim of the NCSP is to detect and treat cervical abnormalities prior to progression to invasive cervical cancer. A national system of state-based Pap Smear Registers (PSR) systematically records results of cervical screening, performs a recall and reminder function for women and their primary care providers, and a safety net function for follow up of screen-detected abnormalities. PSRs provide data for national reporting on screening participation, prevalence of abnormalities, outcomes after an abnormal Pap smear and other program quality indicators [2].

Low-grade abnormalities (LGA), indicated by a cytological diagnosis of a low-grade squamous intraepithelial lesion on a Pap smear, are common among young women and usually represent a cellular response to acute HPV infection [3]. Most LGAs resolve without treatment because the infection is cleared; a repeat smear after 6–12 months is currently recommended (unless a woman is aged over 30 years without a recent normal screening history), because persistent infection increases the risk of a high-grade abnormality being present [4–6]. High-grade abnormalities (HGA), as indicated by a cytological detection of a high-grade squamous intraepithelial lesion or a glandular abnormality, can indicate a true precancerous abnormality of the cervix (cervical intraepithelial neoplasia [CIN] grade 3 or adenocarcinoma in situ) or possibly a CIN grade 2 lesion, which is a diagnostic classification that includes both florid low-grade disease and grade 3 disease [7]. Since it is not possible to determine which high-grade lesions will eventually progress to cancer and which will resolve, all are investigated and treatment of confirmed high grade lesions is recommended by the current National Health and Medical Research Council guidelines (p.53) [4,8]. Prompt follow-up of cytology predicted HGAs by colposcopy and biopsy is required to confirm the diagnosis prior to treating the lesion [4].

Aboriginal and Torres Strait Islander women, hereafter respectfully referred to as Indigenous Australian women, have cervical cancer incidence and mortality rates two and four times higher than their non-Indigenous counterparts [9,10]. The NCSP cannot report on screening participation, abnormalities or outcomes for Indigenous women because pathology report forms (the source of information for PSRs) do not include Indigenous status [10–12].

Recently, using record-linkage methods [13], we have reported lower cervical screening participation rates among Indigenous women compared to non-Indigenous women in Queensland [14]. We report for the first time the prevalence of cervical abnormalities for Indigenous women compared to non-Indigenous women participating in cervical screening in Queensland, where 28.5% of the total Australian Indigenous population residing in this Australian state.

Methods

Data Sources

The detailed data extraction and linkage methods have been described previously [13]. Briefly, the dataset included linked records from the Queensland PSR and the Queensland Hospital

Admitted Patient Data Collections (QHAPDC). The Queensland PSR was used to identify women resident in Queensland aged 20–69 years (at the time of testing) who had a Pap smear between February 1999 (the start of the Queensland PSR) and December 2011, and had not opted to be excluded from the Queensland PSR. Variables obtained from the Queensland PSR included: date of birth (mm/yyyy); place of residence (suburb and postcode); test date; type (cytology or histology); and result. An Indigenous identifier was assigned to the PSR cohort by linking to a QHAPDC extract of women aged 20–69 years who had ever been identified as Indigenous when admitted to a Queensland public hospital during 1995–2011. The QHAPDC has reasonably high accuracy of Indigenous status: in 2011/12, 87% (95% CI 84–91) of Indigenous inpatients were correctly identified as Indigenous in hospital records (compared to self-identification) [15]. Women who linked to at least one QHAPDC record were identified as Indigenous if at least 50% of their QHAPDC records identified them as such [13,16]. Those who did not match to at least one QHAPDC extract record, or had fewer than 50% of their QHAPDC records identified as Indigenous, were assumed to be non-Indigenous. Women were excluded from the study if they had insufficient details to determine the statistical local area (SLA) of residence within Queensland for at least one Pap smear.

Outcome Measures

Abnormal Pap smear results were categorised according to the Australian Modified Bethesda System 2004 as: low-grade abnormality (LGA: possible or definitive low-grade squamous intraepithelial lesion detected at cytology [includes previous terminology of atypical squamous cells of undetermined significance]); or high-grade abnormality (HGA: prediction of CIN 2 or higher, adenocarcinoma *in situ*, or invasive cancers [includes previous terminology of atypical squamous cells, possible high-grade lesion]), consistent with current national reporting [4]. Cytology-detected LGA and HGA are hereafter referred to as cLGA and cHGA, respectively, to distinguish them from histologically confirmed HGA (hHGA). We categorised women as having an hHGA if there was a record of hHGA within six months after the date of a Pap smear. Women with an HGA histology report who had not had a cytology test within the previous six months were excluded because these tests may have resulted from investigations other than cervical screening.

Location of residence based on suburb and postcode at the time of Pap smear was mapped to the 2011 SLA boundaries. If the address information for a specific Pap smear was insufficient to determine SLA, then information from the closest adjacent record for the same woman was used. SLAs were grouped into five categories from major city to very remote [17]. We assigned an area-based measure of socioeconomic disadvantage to each woman based on the SLA of place of residence using the Index of Relative Socio-Economic Advantage and Disadvantage, with Queensland population-based quintiles from most disadvantaged to most affluent [18].

Statistical analyses. Demographic characteristics are presented as medians (with interquartile ranges [IQR]) for non-normally distributed continuous variables and as frequencies and percentages for categorical variables. Proportions for different groups of women were compared using chi-squared tests. Prevalence of hHGA was determined by dividing the number of hHGA in each two-year calendar period (2000–2001 to 2010–2011) by the number of women who were screened in the corresponding two-year period and directly age-standardised based on the 2001 Australian Estimated Resident Population and expressed per 1000 women [19]. Simple linear regression was used to graphically present the association of age-standardised hHGA prevalence and time period stratified by Indigenous status.

Logistic regression was used to quantify the association (as odds ratios) between independent variables of interest and the prevalence of each of cLGA, cHGA and hHGA. The regression

models for each outcome included five a priori independent categorical variables of interest (the 'main effects' model): Indigenous status; age at time of test, in age-groups of 20–29, 30–39, 40–49 (reference category), 50–59 and 60–69 years; two-year calendar periods from 2000–2001 (reference category) to 2010–2011; place of residence (inner regional as the reference category); and disadvantage quintiles (quintile 3 as the reference category). An interaction between each independent variable and Indigenous status was assessed, but these were not included in the final models because inclusion of the individual interaction terms did not change the estimates for the other a priori variables substantially. To examine if prevalence of abnormalities within each age-group was changing over time, we fitted separate logistic models for each age-group for cLGA and hHGA, including two-year calendar periods as an ordinal variable and Indigenous status, area-level disadvantage and place of residence. We assessed if temporal trends differed by Indigenous status by fitting an interaction term.

To assess whether hHGA was more common for women who had not previously had a Pap smear, we selected women aged 30–69 years who had a Pap smear in 2010–2011 and dichotomised their screening history into 'yes previous screen' or 'no previous screen' using a ten-year look-back period (2000–2009). Of these women, we determined how many had an hHGA up to six months after their Pap smear in 2010–2011.

Given the non-independence of multiple tests carried out for each woman, we accounted for clustering (i.e. treating each woman as a cluster) in the logistic models and assessed the overall model fit. Because the likelihood ratio test was not appropriate when accounting for clustering, we used the Wald chi-squared test to determine overall significance of each covariate and interaction term. Joint chi-squared tests were used to assess the contribution of each variable to model fit, and the Z test to assess the significance of individual coefficients within the logistic model.

All analyses were conducted using Stata (Version 14.0; Stata Corporation, College Station, TX) [20].

The Human Research Ethics Committees (HREC) of Queensland Health (Far North Queensland HREC HREC/15/QCH/19-957), the Northern Territory Department of Health & Menzies School of Health Research (HOMER-2012-1737) and Charles Darwin University (H12093) approved the study along with the Queensland Research Linkage Group, data custodians, and director general of Queensland Health to access and link records. The research team received a de-identified linked dataset and therefore was unable to obtain individual consent from participants. This process was approved by the aforementioned ethics committees.

Results

We excluded 1545 women with conflicting dates of birth, 3174 women with missing address details (a total of 11,072 Pap tests) and 518 women with hHGA only. Our final cohort included 1,334,795 women with 4,565,250 Pap smears from 2000–2011. There were 26,829 Indigenous women (2.0%) identified in the PSR cohort with 87,372 Pap smears. The median number of Pap smears per woman was similar for Indigenous and non-Indigenous women (3, IQR 1–5). Similar proportions of Indigenous and non-Indigenous women had only one (26.9% vs 26.6%), two or three (35.4% vs 30.9%) and four or more (37.7% vs 42.4%) Pap smears during the study period. The demographic details of women at their first recorded Pap smear are summarised in [Table 1](#). Indigenous women compared to non-Indigenous women were more likely to be younger (median age 34, IQR 27–43 vs 40, IQR 30–50), live outside major cities and live in less affluent areas.

The proportions of abnormal tests by socio-demographic variables for Indigenous and non-Indigenous women are shown in [Table 2](#). Greater proportions of tests for Indigenous women

Table 1. Characteristics of Indigenous and non-Indigenous women at their first recorded Pap smear.

Variables	Indigenous		Non-Indigenous		p ^a
	N	%	N	%	
Totals	26,829	2.0	1,307,966	98.0	
Age group (years)					
20–29	13,104	48.8	476,371	36.4	
30–39	6790	25.3	388,932	25.9	
40–49	4107	15.3	258,661	19.8	
50–59	1982	7.4	158,310	12.1	
60–69	846	3.2	75,692	5.8	<0.001
Area-level disadvantage					
Q1 (most disadvantaged)	7358	27.4	148,552	11.4	
Q2	9281	34.6	285,133	21.8	
Q3	4877	18.2	301,630	23.1	
Q4	4090	15.2	326,672	25.0	
Q5 (most affluent)	1223	4.6	245,979	18.8	<0.001
Place of residence					
Major city	6700	25.0	812,894	62.2	
Inner regional	4259	15.9	252,875	19.3	
Outer regional	10,593	39.5	218,651	16.7	
Remote	2129	7.9	14,611	1.1	
Very remote	3148	11.7	8935	0.7	<0.001

^aChi-squared test of association for categorical variables

doi:10.1371/journal.pone.0150473.t001

compared with non-Indigenous women had cLGA and cHGA (p<0.001) overall and this was consistent by age-group, area-level disadvantage and place of residence.

The odds of Indigenous women having cLGA (OR 1.7, 95% CI 1.6–1.7), cHGA (OR 2.8, CI 2.7–3.0) and hHGA (OR 2.6, CI 2.5–2.8) were higher than non-Indigenous women after adjusting only for year of Pap smear. The odds remained higher when adjusted further for area-level disadvantage, place of residence and age-group, for example, cLGA OR 1.4, CI 1.3–1.4 (Table 3).

The prevalence of cLGA decreased after 2005; was lower for older than younger women, but was not associated with area-level disadvantage. Compared to women in inner regional areas, women living in all other areas had a higher prevalence of cLGA. For all women, cLGA prevalence significantly decreased over time in all age-groups (Table 4). The prevalence of cHGA increased over time, was lower in the most affluent area, and for older women, but was higher in outer regional, remote and very remote areas (Table 3).

The prevalence of hHGA was higher among Indigenous (16.5 per 1000 women screened, CI 13.9–19.5) than non-Indigenous women (7.3 per 1000 women screened, CI 7.1–7.6), in 2000–2001 and remained relatively constant in both groups up to 2010–2011 (16.6 per 1000 women screened, CI 14.6–18.9 and 7.5 per 1000 women screened, CI 7.3–7.7, respectively). Prevalence was higher for Indigenous women in each age-group (Fig 1). In the 20–29 year age-group, prevalence of hHGA increased over time for Indigenous women only and multivariable analysis confirmed this was a statistically significant trend (Table 4). There were no significant changes in prevalence of hHGA over time for other age-groups among Indigenous women. Among non-Indigenous women, hHGA prevalence decreased significantly in 40–69 year olds.

Table 2. Demographic characteristics of Indigenous and non-Indigenous women’s Pap smears by abnormality, 2000–2011.

Variables	Indigenous			Non-Indigenous		
	N	% cLGA	% cHGA	N	% cLGA	% cHGA ^a
No. Pap smears	87,372	6.6	3.2	4,477,878	4.3	1.2
Age group (years)						
20–24	15,485	13.4	5.7	481,149	10.9	2.3
25–29	15,897	8.5	5.2	554,668	6.8	2.2
30–39	26,854	6.0	3.1	1,193,327	4.3	1.4
40–49	16,659	4.7	2.1	1,080,806	3.7	0.7
50–59	8619	3.4	1.3	765,180	2.8	0.5
60–69	3 858	3.4	1.5	402,748	2.0	0.5
Area-level disadvantage						
Q1	24,546	7.2	3.6	501,561	4.5	1.2
Q2	30,114	7.0	3.7	996,147	4.6	1.3
Q3	15,363	7.2	3.3	1,042,366	4.6	1.2
Q4	13,164	7.4	3.6	1,080,243	4.8	1.2
Q5 (most affluent)	4185	6.9	2.7	857,561	4.8	1.1
Place of residence						
Major cities	21,334	7.0	3.1	2,745,136	4.8	1.1
Inner regional	13,503	7.1	3.4	895,540	4.2	1.2
Outer regional	35,060	7.0	3.7	761,111	4.9	1.3
Remote	6609	8.4	4.4	46,932	5.0	1.6
Very remote	10,866	7.0	3.3	29,159	4.9	1.7

- Notes:
- Table 2 shows the proportion of abnormal results from the total number of tests by Indigenous status (excluding tests that were taken in the same year with the same result).
 - cLGA: Cytology detected low-grade abnormality; cHGA: cytology detected high-grade abnormality.
- ^a Statistically significant chi-squared test for association, $p < 0.001$ comparing the proportion of abnormal Pap smears among Indigenous women and the proportion of abnormal Pap smears among non-Indigenous women.

doi:10.1371/journal.pone.0150473.t002

In 2010–2011, 548,549 women aged 30–69 years had a Pap smear, of whom 1.6% were Indigenous. hHGA was more common among women without a Pap smear history (unadjusted prevalence Indigenous: 36.6 per 1000 women without and 11.3 per 1000 with screening history; non-Indigenous: 10.7 per 1000 without and 4.5 per 1000 with screening history).

We carried out a sensitivity analysis by re-running all models and estimating hHGA prevalence by excluding cervical cancer in the definition of the outcome variable; there were no differences in these and the main results for the prevalence of hHGA (comparison data not shown). We also calculated a histology detection rate, based on the number of HGAs reported by histology, using the method as reported by the Australian Institute of Health and Welfare [21]: the detection rate was 23 and 9 per 1000 women screened in 2011 for Indigenous women and non-Indigenous women, respectively.

Discussion

Indigenous women had markedly higher prevalence of cytology-detected and histology-confirmed cervical abnormalities than non-Indigenous women. The prevalence of cHGA increased over time for Indigenous women only. Prevalence of cLGA decreased over time for both groups of women.

Table 3. Association of Indigenous status, calendar period, place of residence, area-level disadvantage and age with risk of cytology detected low-grade abnormalities and cytological and histological high-grade abnormalities including cervical cancer.

Variables	cLGA Adjusted ORs (95% CI)	cHGA Adjusted ORs (95% CI)	hHGA Adjusted ORs (95% CI)
Indigenous status			
Indigenous women	1.38 (1.33–1.43)	2.16 (2.05–2.27)	1.98 (1.87–2.10)
Year			
2000–2001	1.00	1.00	1.00
2002–2003	0.99 (0.98–1.01)	1.04 (1.00–1.07)	1.07 (1.03–1.11)
2004–2005	0.99 (0.97–1.00)	1.19 (1.15–1.23)	0.98 (0.94–1.01)
2006–2007	0.82 (0.80–0.83)	1.23 (1.19–1.287)	0.92 (0.88–0.96)
2008–2009	0.65 (0.64–0.66)	1.19 (1.15–1.23)	0.94 (0.91–0.98)
2010–2011	0.52 (0.57–0.59)	1.25 (1.25–1.33)	1.02 (0.98–1.06)
Area-level disadvantage			
Q1	1.00 (0.98–1.02)	1.06 (1.02–1.10)	1.04 (1.00–1.09)
Q2	1.01 (0.99–1.03)	1.06 (1.03–1.09)	1.08 (1.04–1.12)
Q3	1.00	1.00	1.00
Q4	1.01 (0.99–1.03)	1.02 (0.99–1.05)	0.97 (0.94–1.00)
Q5 (most affluent)	1.00 (0.98–1.02)	0.90 (0.87–0.93)	0.94 (0.88–1.02)
Place of residence			
Major city	1.09 (1.08–1.11)	0.99 (0.96–1.02)	1.02 (0.98–1.05)
Inner regional	1.00	1.00	1.00
Outer regional	1.12 (1.10–1.14)	1.09 (1.05–1.12)	1.07 (1.03–1.11)
Remote	1.13 (1.08–1.18)	1.24 (1.15–1.34)	1.18 (1.08–1.30)
Very remote	1.09 (1.04–1.16)	1.11 (1.02–1.22)	1.04(0.93–1.16)
Age group (years)			
20–29	2.53 (2.50–2.57)	3.15 (2.05–2.27)	4.20 (4.05–4.36)
30–39	1.18 (1.16–1.20)	1.87 (1.81–1.92)	2.30 (2.21–2.39)
40–49	1.00	1.00	1.00
50–59	0.75 (0.74–0.77)	0.70 (0.67–0.73)	0.53 (0.50–0.56)
60–69	0.52 (0.50–0.53)	0.59 (0.56–0.63)	0.42 (0.39–0.46)

Notes:

1. Binary logistic regression model comparing odds of having an abnormality adjusted for all a priori variables in the table. Clustering of Pap smears for women was accounted for in each model.
2. Table 3 presents OR, Odd Ratios; 95% C.I., 95% Confidence Intervals
3. Each independent variable in the table had overall $p < 0.001$ and each outcome variable (type of abnormality) $p < 0.001$.

doi:10.1371/journal.pone.0150473.t003

We were able to identify Indigenous women in the PSR only if women had been a public hospital inpatient between 1995 and 2011 [13]. Women admitted to hospital may differ from those who were not. Indirect evidence suggests that almost all Indigenous women in the age-range for our study would have been admitted to a public hospital over the 17 year period that we used to identify Indigenous women [13], so it is likely that our results apply to almost all screening women. Queensland has the second highest proportion of Indigenous people in its population and they have widely varied geographic and socioeconomic characteristics. Therefore, it is probable that the results for Queensland be applicable to the broader Australian Indigenous population.

The availability of individual-level PSR data facilitated a more precise method for estimating the prevalence of hHGA than reported previously [21]. This previous method calculated hHGA

Table 4. Time trends in prevalence of cytology low-grade and histologically confirmed high-grade abnormalities by age-group, 2000–2011.

Age-group ^a	cLGA	hHGA	
	All women ^b	Indigenous	Non-Indigenous
20–29	0.94 (0.94–0.94)	1.07 (1.03–1.12)	1.00 (0.99–1.01)
30–39	0.89 (0.88–0.85)	0.99 (0.93–1.05)	0.99 (0.98–1.01)
40–49	0.84 (0.84–0.85)	1.03 (0.93–1.14)	0.96 (0.94–0.98)
50–59	0.81 (0.81–0.82)	0.92 (0.76–1.12)	0.94 (0.91–0.97)
60–69	0.81 (0.80–0.82)	0.84 (0.68–1.02)	0.91 (0.87–0.96)

Notes:

1. Table 4 reports OR (Odd Ratios) for a two-year increase in time; followed by 95% CI (Confidence Intervals).

2. cLGA: Cytology detected low-grade abnormality; cHGA: cytology detected high-grade abnormality.

^aeach model included Indigenous status, area-level disadvantage and place of residence and an interaction term for Indigenous status and year.

^bORs were similar for Indigenous and non-Indigenous women for cLGAs over time and therefore have been presented for all women combined.

doi:10.1371/journal.pone.0150473.t004

prevalence from aggregated data supplied by PSRs: number of hHGAs (excluding cancer) divided by the number of women screened. We calculated hHGA prevalence directly for screened women (i.e. including only hHGA/cervical cancer if there was a record of a recent Pap smear) using individual-level data. This increased precision in estimating hHGA prevalence in screened women provides greater validity of our study findings. Differences in estimation methods mean comparisons with previously published estimates requires some caution [22].

Participation in cervical screening of Indigenous women is significantly less than that of non-Indigenous women in Queensland, a known risk factor for cervical cancer [14]. Increasing screening among Indigenous women would likely reduce HGAs. We did not have data about other risk factors such as persistent HPV infection, lower individual-level socio-economic status, lower education level, smoking, possible dietary deficiencies, oral contraception, age at first intercourse and early and higher parity [23–27]. The prevalence of HPV types 16 and 18 has previously been reported to be similar for Indigenous and non-Indigenous women and more recently similar age at sexual debut has been reported [28,29]. Some evidence indicates that Indigenous women have higher prevalence of other oncogenic HPV types in the 36–40 year age-group [29]. Indigenous women in this cohort had much higher prevalence of LGAs—a known marker of productive HPV infection—and hHGAs, of which 50–60% are caused by HPV types 16 and 18 [3,30]. This suggests that a difference in HPV infection patterns may exist. Young Indigenous women have higher smoking rates [2], lower age at parity and higher fertility rates [31], less oral contraceptive use [29] and are more likely to live in areas of disadvantage [2]. It is crucial to understand the possible role of these factors in the increased risk of HGAs developing (or not regressing) among Indigenous women.

The Australian Government has announced changes to the current NCSP which will shift to a new program (the ‘Renewal’) in May 2017 using primary HPV testing every five years for women aged 25–74 years [32,33]. Starting screening at age 25 is consistent with international reports showing that screening and treatment of HGAs below the age of 25 is not effective at preventing cancer [34], and will occur in the context of the significant decline in high-grade lesions in 20–24 year olds following the 2007 introduction of the national HPV vaccination program in Australia [10,35]. Given our findings it is timely and important that the Renewal helps address the needs of Indigenous women. High prevalence of HGAs among Indigenous women has previously been reported among Indigenous women in remote areas [22]. Increasing the uptake of HPV vaccination for Indigenous children must be a priority. Although some evidence indicates coverage and/or dose completion rates for Indigenous female adolescents




Fig 1. Prevalence of histologically confirmed high-grade abnormalities per 1000 women screened, by age-group and Indigenous status, 2000–2011.

doi:10.1371/journal.pone.0150473.g001

during the catch-up phase were lower than for non-Indigenous female adolescents [36], reassuringly, early vaccination program outcomes in terms of falls in presentations for anogenital warts (caused by HPV types also included in the vaccine) have been similar for 15–25 year old Indigenous and non-Indigenous females [37]. Over the longer term, it is hoped that current school-based routine HPV vaccination, which now includes boys as well as girls, will encourage relatively high ongoing coverage overall and that this mode of delivery will reduce inequities in outcomes for Indigenous women.

The Renewal will provide the option of self-collection of HPV samples for under-screened and never-screened women, which is supported by international evidence that self-collection can increase screening among these groups [38]. Other opportunities to increase screening include an invitation to commence screening at a woman's 25th birthday using a call-and-recall system inviting women to rescreen, rather than the current reminder-based screening system. Possible mediums of invitations could be considered such as using SMS or email or letter and the inclusion of educational resources and information.

Evidence suggests that there has been some reduction in the incidence of cervical cancer over time among Indigenous women [9,10,39], indicating that efforts to reduce the burden of cervical cancer for Indigenous women have been somewhat effective. For over two decades the NCSP has been unable to report on the impacts of the program for Indigenous Australian women on a wide range of outcomes [11,12,21]. The Renewal program must take the critical opportunity to ensure that Indigenous women are able to be recorded in the proposed national registry. Our study provides benchmark measures giving an opportunity to develop and assess the effectiveness of interventions designed to increase the number of women who are followed up for a serious abnormality and reduce the severity of those outcomes.

Internationally, Australia has been at the forefront of cervical cancer prevention. However, below the surface of this success, Indigenous Australian women carry a disproportionate burden of disease, comparable to many countries without the resources available in Australia. The Renewal of the cervical screening program provides the opportunity to overcome long standing data deficiencies and should consider the Indigenous-specific patterns and trends identified in our study. It is imperative that any major changes to current screening practices prioritise Indigenous women. Failure to do so will result in further disenfranchisement and disadvantage of Indigenous women, their families and their communities.

Acknowledgments

We would like to acknowledge the staff and registrars from the Queensland Pap Smear Register, the Queensland Cancer Registry and the Queensland Health Admitted Patient Data Collections for their assistance in providing the data, and the Queensland Research Linkage-Group for linking the data. We gratefully acknowledge Tegan Harris for a range of contributions, including building and designing the database.

Author Contributions

Conceived and designed the experiments: LJW JRC PB GG JC PCV DO JMLB KC KL DR DG. Analyzed the data: LJW. Wrote the paper: LJW JRC PB GG JC PCV DO JMLB KC KL DR DG SPM AD. Contributed critically to the interpretation of study findings: LJW JRC PB GG JC PCV DO JMLB KC KL DR DG SPM AD. Conducted the approval process and data acquisition: LJW GG JRC AD SPM. Provided data analysis support: PB JRC JC AD KL.

References

1. Australian Government Department of Health (2013) National Cervical Screening Program. Canberra.

2. Australian Institute of Health and Welfare (2015) The health and welfare of Australia's Aboriginal and Torres Strait Islander peoples 2015. Cat. no. IHW 147. Canberra: AIHW.
3. Schiffman M, Wentzensen N (2013) Human papillomavirus infection and the multistage carcinogenesis of cervical cancer. *Cancer Epidemiol Biomarkers Prev* 22: 553–560. doi: [10.1158/1055-9965.EPI-12-1406](https://doi.org/10.1158/1055-9965.EPI-12-1406) PMID: [23549399](https://pubmed.ncbi.nlm.nih.gov/23549399/)
4. National Health and Medical Research Council (2005) Screening to Prevent Cervical Cancer: Guidelines for the Management of Asymptomatic Women with Screen Detected Abnormalities. Canberra: NHMRC.
5. Carter JR, Ding Z, Rose BR (2011) HPV infection and cervical disease: a review. *Aust N Z J Obstet Gynaecol* 51: 103–108. doi: [10.1111/j.1479-828X.2010.01269.x](https://doi.org/10.1111/j.1479-828X.2010.01269.x) PMID: [21466509](https://pubmed.ncbi.nlm.nih.gov/21466509/)
6. Sellors JW, Sankaranarayanan R (2003) Chapter 2: An introduction to cervical intraepithelial neoplasia (CIN). *Colposcopy and treatment of cervical intraepithelial neoplasia: A beginner's manual*. Lyon, France: International Agency for Research on Cancer.
7. Schiffman M, Castle PE, Jeronimo J, Rodriguez AC, Wacholder S (2007) Human papillomavirus and cervical cancer. *Lancet* 370: 890–907. PMID: [17826171](https://pubmed.ncbi.nlm.nih.gov/17826171/)
8. McCredie MR, Sharples KJ, Paul C, Baranyai J, Medley G, Jones RW, et al. (2008) Natural history of cervical neoplasia and risk of invasive cancer in women with cervical intraepithelial neoplasia 3: a retrospective cohort study. *Lancet Oncol* 9: 425–434. doi: [10.1016/S1470-2045\(08\)70103-7](https://doi.org/10.1016/S1470-2045(08)70103-7) PMID: [18407790](https://pubmed.ncbi.nlm.nih.gov/18407790/)
9. Zhang X, Condon JR, Rumbold AR, Cunningham J, Roder DM (2011) Estimating cancer incidence in Indigenous Australians. *Aust N Z J Public Health* 35: 477–485. doi: [10.1111/j.1753-6405.2011.00762.x](https://doi.org/10.1111/j.1753-6405.2011.00762.x) PMID: [21973255](https://pubmed.ncbi.nlm.nih.gov/21973255/)
10. Australian Institute of Health and Welfare (2015) Cervical screening in Australia 2012–2013. Cancer series 93. Cat. no. CAN 91. Canberra: AIHW.
11. Australian Institute of Health and Welfare (2013) The inclusion of Indigenous status on pathology request forms. Cat. no. IHW 103 Canberra: AIHW.
12. Whop LJ, Cunningham J, Condon JR (2014) How well is the National Cervical Screening Program performing for Indigenous Australian women? Why we don't really know, and what we can and should do about it. *Eur J Cancer Care (Engl)* 23: 716–720.
13. Whop LJ, Diaz A, Baade P, Garvey G, Cunningham J, Brotherton JM, et al. (2016) Using probabilistic record linkage methods to identify Australian Indigenous women on the Queensland Pap Smear Register: the National Indigenous Cervical Screening Project. *BMJ Open* 6: e009540. doi: [10.1136/bmjopen-2015-009540](https://doi.org/10.1136/bmjopen-2015-009540) PMID: [26873047](https://pubmed.ncbi.nlm.nih.gov/26873047/)
14. Whop LJ, Garvey G, Baade P, Lokuge K, Cunningham J, Brotherton JML, et al. (in press (2016)) The first comprehensive report on Indigenous Australian women's inequalities in cervical screening: a retrospective registry cohort study in Queensland, Australia (2000–2011). *Cancer*.
15. Australian Institute of Health and Welfare (2013) Indigenous identification in hospital separations data-Quality report. Cat. no. IHW 90. Canberra: AIHW.
16. Australian Institute of Health and Welfare and Australian Bureau of Statistics (2012) National best practice guidelines for data linkage activities relating to Aboriginal and Torres Strait Islander people. Canberra: AIHW.
17. Australian Institute of Health and Welfare (2004) Rural, Regional and Remote Health, A Guide to Remoteness Classifications. Canberra: AIHW.
18. Australian Bureau of Statistics (2011) Technical Paper Socio-Economic Indexes for Areas (SEIFA) 2011. Canberra: ABS.
19. Australian Institute of Health and Welfare (2011) Principles on the use of direct age-standardisation in administrative data collections: for measuring the gap between Indigenous and non-Indigenous Australians. Canberra: AIHW.
20. StataCorp. (2015) Stata Statistical Software: Release 14. College Station, TX: StataCorp LP.
21. Australian Institute of Health and Welfare (2014) Cervical screening in Australia 2011–2012. Cancer series 82. Cat. no. CAN 72. Canberra: AIHW.
22. Zhang X, Condon JR, Douglas F, Bates D, Gutheridge S, Garling L, et al. (2012) Women's cancers and cancer screening in the Northern Territory. Darwin: Department of Health.
23. Jensen KE, Schmiedel S, Norrild B, Frederiksen K, Iftner T, Kjaer SK (2013) Parity as a cofactor for high-grade cervical disease among women with persistent human papillomavirus infection: a 13-year follow-up. *Br J Cancer* 108: 234–239. doi: [10.1038/bjc.2012.513](https://doi.org/10.1038/bjc.2012.513) PMID: [23169283](https://pubmed.ncbi.nlm.nih.gov/23169283/)
24. Franceschi S, Plummer M, Clifford G, de Sanjose S, Bosch X, Herrero R, et al. (2009) Differences in the risk of cervical cancer and human papillomavirus infection by education level. *Br J Cancer* 101: 865–870. doi: [10.1038/sj.bjc.6605224](https://doi.org/10.1038/sj.bjc.6605224) PMID: [19654578](https://pubmed.ncbi.nlm.nih.gov/19654578/)

25. International Collaboration of Epidemiological Studies of Cervical Cancer (2007) Cervical cancer and hormonal contraceptives: collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women without cervical cancer from 24 epidemiological studies. *The Lancet* 370: 1609–1621.
26. Smith JS, Green J, Berrington de Gonzalez A, Appleby P, Peto J, Plummer M, et al. (2003) Cervical cancer and use of hormonal contraceptives: a systematic review. *Lancet* 361: 1159–1167. PMID: [12686037](#)
27. International Collaboration of Epidemiological Studies of Cervical C, Appleby P, Beral V, Berrington de Gonzalez A, Colin D, Franceschi S, et al. (2006) Carcinoma of the cervix and tobacco smoking: collaborative reanalysis of individual data on 13,541 women with carcinoma of the cervix and 23,017 women without carcinoma of the cervix from 23 epidemiological studies. *Int J Cancer* 118: 1481–1495. PMID: [16206285](#)
28. Ward J, J. Bryant, Wand H, Pitts M, Smith A, Delaney-Thiele D, et al. (2014) Sexual Health and relationships in young Aboriginal and Torres Strait Islander people: Results from the first national study assessing knowledge, risk practices and health service use in relation to sexually transmitted infections and blood borne viruses (The Goanna Survey).
29. Garland SM, Brotherton JM, Condon JR, McIntyre PB, Stevens MP, Smith DW, et al. (2011) Human papillomavirus prevalence among indigenous and non-indigenous Australian women prior to a national HPV vaccination program. *BMC Med* 9: 104. doi: [10.1186/1741-7015-9-104](#) PMID: [21910918](#)
30. Wright TC Jr. (2006) HPV DNA testing for cervical cancer screening. FIGO 26th Annual Report on the Results of Treatment in Gynecological Cancer. *Int J Gynaecol Obstet* 95 Suppl 1: S239–246. PMID: [17161163](#)
31. Australian Bureau of Statistics (2011) Patient Experiences in Australia: Summary of Findings, 2012–13. Cat No 4839.0. Canberra: ABS.
32. MSAC (Medical Services Advisory Committee) (2014) MSAC application no.1276: National Cervical Screening Program renewal. Canberra: MSAC.
33. Australian Government Department of Health (2015) Overview of the Renewal. Canberra.
34. Sasieni P, Castanon A, Cuzick J (2009) Effectiveness of cervical screening with age: population based case-control study of prospectively recorded data. *BMJ* 339: b2968. doi: [10.1136/bmj.b2968](#) PMID: [19638651](#)
35. Brotherton JM, Saville AM, May CL, Chappell G, Gertig DM (2015) Human papillomavirus vaccination is changing the epidemiology of high-grade cervical lesions in Australia. *Cancer Causes Control* 26: 953–954. doi: [10.1007/s10552-015-0568-6](#) PMID: [25804857](#)
36. Brotherton JM, Murray SL, Hall MA, Andrewartha LK, Banks CA, Meijer D, et al. (2013) Human papillomavirus vaccine coverage among female Australian adolescents: success of the school-based approach. *Med J Aust* 199: 614–617. PMID: [24182228](#)
37. Smith MA, Liu B, McIntyre P, Menzies R, Dey A, Canfell K (2015) Fall in genital warts diagnoses in the general and indigenous Australian population following implementation of a national human papillomavirus vaccination program: analysis of routinely collected national hospital data. *J Infect Dis* 211: 91–99. doi: [10.1093/infdis/jiu370](#) PMID: [25117753](#)
38. Arbyn M, Verdoodt F, Snijders PJ, Verhoef VM, Suonio E, Dillner L, et al. (2014) Accuracy of human papillomavirus testing on self-collected versus clinician-collected samples: a meta-analysis. *Lancet Oncol* 15: 172–183. doi: [10.1016/S1470-2045\(13\)70570-9](#) PMID: [24433684](#)
39. Zhang X C J, Dempsey K, Garling L., (2014) Cancer in the Northern Territory 1991–2010: incidence, mortality and survival. Darwin: Department of Health.