

---

Charles Darwin University

## Complete Genome Sequence of the Encephalomyelitic *Burkholderia pseudomallei* Strain MSHR305

Stone, Joshua; Johnson, Shannon; Bruce, David; Detter, John Chris; Mayo, Mark; Currie, Bart; Gelhaus, H. Carl; Keim, Paul S; Tuanyok, Apichai

*Published in:*  
Genome Announcements (genomeA)

*DOI:*  
[10.1128/genomeA.00656-13](https://doi.org/10.1128/genomeA.00656-13)

Published: 01/01/2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

### *Citation for published version (APA):*

Stone, J., Johnson, S., Bruce, D., Detter, J. C., Mayo, M., Currie, B., Gelhaus, H. C., Keim, P. S., & Tuanyok, A. (2013). Complete Genome Sequence of the Encephalomyelitic *Burkholderia pseudomallei* Strain MSHR305. *Genome Announcements (genomeA)*, 1(4), 1-2. [e00656-13]. <https://doi.org/10.1128/genomeA.00656-13>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Complete Genome Sequence of the Encephalomyelitic *Burkholderia pseudomallei* Strain MSHR305

Joshua K. Stone,<sup>a</sup> Shannon L. Johnson,<sup>b</sup> David C. Bruce,<sup>b</sup> J. Chris Detter,<sup>b</sup> Mark Mayo,<sup>c</sup> Bart J. Currie,<sup>c</sup> H. Carl Gelhaus,<sup>d</sup> Paul Keim,<sup>a</sup> Apichai Tuanyok<sup>a</sup>

Department of Biological Sciences, Northern Arizona University, Flagstaff, Arizona, USA<sup>a</sup>; Los Alamos National Laboratory, Los Alamos, New Mexico, USA<sup>b</sup>; Menzies School of Health Research, Charles Darwin University, Darwin, Northern Territory, Australia<sup>c</sup>; Battelle Memorial Institute, Columbus, Ohio, USA<sup>d</sup>

**We describe the complete genome sequence of *Burkholderia pseudomallei* MSHR305, a clinical isolate taken from a fatal encephalomyelitis case, a rare form of melioidosis. This sequence will be used for comparisons to identify the genes that are involved in neurological cases.**

Received 23 July 2013 Accepted 24 July 2013 Published 22 August 2013

**Citation** Stone JK, Johnson SL, Bruce DC, Detter JC, Mayo M, Currie BJ, Gelhaus HC, Keim P, Tuanyok A. 2013. Complete genome sequence of the encephalomyelitic *Burkholderia pseudomallei* strain MSHR305. *Genome Announc.* 1(4):e00656-13. doi:10.1128/genomeA.00656-13.

**Copyright** © 2013 Stone et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Apichai Tuanyok, Apichai.Tuanyok@nau.edu.

*Burkholderia pseudomallei* is a Gram-negative saprophytic bacterium endemic to southeast Asia and northern Australia (1). It is the etiological agent of melioidosis, a disease that can present in numerous forms, from cutaneous to pneumonia to septicemia (2). Neurological cases occur rarely, in only 3 to 4% of patients, and in murine models, inhalation led to direct brain infection, presumably through the nasal mucosa via the olfactory nerve (2–4). *B. pseudomallei* strain MSHR305 was isolated in 1994 during an autopsy of a fatal melioidosis encephalomyelitis case at the Royal Darwin Hospital, Northern Territory, Australia, and was previously sequenced to 36 contigs (5).

DNA sequencing was performed by the Los Alamos National Laboratory (LANL) Genome Science Group using Illumina and PacBio technologies (6, 7). Short-insert paired-end Illumina libraries yielded 27,714,112 reads totaling 27.7 Mbp, and long-insert libraries generated 920,819 paired reads totaling 159 Mbp. The PacBio library generated 260,870 subreads for a total of 570 Mbp. The Illumina short-insert and long-insert paired data were assembled together using Newbler version 2.6. The Newbler consensus sequences were computationally shredded into 2-kbp overlapping short reads (shreds). The Illumina data were also assembled with Velvet version 1.2.08 (8), and the consensus sequence was shredded into 1.5-kbp overlapping shards. The draft data from all platforms were assembled together with AllPaths, version 39750, and the consensus sequence was shredded into 10-kbp overlapping shreds. We integrated consensus shreds from Newbler, Velvet, and AllPaths along with a subset of read pairs from the Illumina long-insert library using parallel Phrap version SPS-4.24 (High Performance Software, LLC). Gap closure was accomplished using PacBio consensus sequences.

The final assembly consisted of two chromosomes of lengths 4,054,138 and 3,373,904 bp, with average coverages of 287× and 79.2× for the Illumina and PacBio reads, respectively, and 6,209 total predicted genes. A comparison to the prototypic strain *B. pseudomallei* K96243 and to *B. pseudomallei* MSHR668, an isolate from a nonfatal encephalomyelitis case (5), revealed the pres-

ence of a hemolysin and activator protein on chromosome 1 in MSHR305 and MSHR668 that is not present in K96243. However, BLAST searches reveal that these proteins are in most *B. pseudomallei* strains. With the results of Owen et al. (4), it appears that most strains have the capacity to cause neurological melioidosis if the bacteria are inhaled and travel directly to the brain. We note that MSHR305 is one of the six priority *B. pseudomallei* strains that were selected as challenge materials for medical countermeasure (MCM) efficacy testing by the Biomedical Advanced Research and Development Authority (BARDA) (9).

**Nucleotide sequence accession numbers.** This whole-genome sequencing project has been deposited at GenBank under the accession no. CP006469 and CP006470. The version described in this paper is the first version.

## ACKNOWLEDGMENTS

This research was financially supported by the Commonwealth of Australia through the National Security Science and Technology Centre within the Defence Science and Technology Organisation and by U.S. Department of Homeland Security (DHS) grant no. HSHQDC-10-C-00149 and HSHQDC-10-00135 and BARDA contract no. HHSO10020110000051. These supports do not represent an endorsement of the contents or conclusions of the research.

## REFERENCES

1. Cheng AC, Currie BJ. 2005. Melioidosis: epidemiology, pathophysiology, and management. *Clin. Microbiol. Rev.* 18:383–416.
2. White NJ. 2003. Melioidosis. *Lancet* 361:1715–1722.
3. Currie BJ, Ward L, Cheng AC. 2010. The epidemiology and clinical spectrum of melioidosis: 540 cases from the 20 year Darwin prospective study. *PLoS Negl. Trop. Dis.* 4:e900. doi:10.1371/journal.pntd.0000900.
4. Owen SJ, Batzloff M, Chehrehasa F, Meedeniya A, Casart Y, Logue CA, Hirst RG, Peak IR, Mackay-Sim A, Beacham IR. 2009. Nasal-associated lymphoid tissue and olfactory epithelium as portals of entry for *Burkholderia pseudomallei* in murine melioidosis. *J. Infect. Dis.* 199:1761–1770.
5. Tuanyok A, Leadem BR, Auerbach RK, Beckstrom-Sternberg SM, Beckstrom-Sternberg JS, Mayo M, Wuthiekanun V, Brettin TS, Nierman WC, Peacock SJ, Currie BJ, Wagner DM, Keim P. 2008. Genomic islands

- from five strains of *Burkholderia pseudomallei*. BMC Genomics 9:566. doi: [10.1186/1471-2164-9-566](https://doi.org/10.1186/1471-2164-9-566).
6. Bennett S. 2004. Solexa Ltd. Pharmacogenomics 5:433–438.
  7. Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D, et al. 2009. Real-time DNA sequencing from single polymerase molecules. Science 323:133–138.
  8. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. Genome Res. 18:821–829.
  9. Van Zandt KE, Tuanyok A, Keim PS, Warren RL, Gelhaus HC. 2012. An objective approach for *Burkholderia pseudomallei* strain selection as challenge material for medical countermeasures efficacy testing. Front. Cell. Infect. Microbiol. 2:120. doi:[10.3389/fcimb.2012.00120](https://doi.org/10.3389/fcimb.2012.00120).