
Charles Darwin University

Quantitative PCR confirms culture as the gold standard for detection of lower airway infection by nontypeable *Haemophilus influenzae* in Australian Indigenous children with bronchiectasis

Hare, Kim; Marsh, Robyn; Binks, Michael; Grimwood, Keith; Pizzutto, Susan; Leach, Amanda; Chang, Anne; Smith-Vaughan, Heidi

Published in:
Journal of Microbiological Methods

DOI:
[10.1016/j.mimet.2012.12.013](https://doi.org/10.1016/j.mimet.2012.12.013)

Published: 01/03/2013

Document Version
Early version, also known as pre-print

[Link to publication](#)

Citation for published version (APA):

Hare, K., Marsh, R., Binks, M., Grimwood, K., Pizzutto, S., Leach, A., Chang, A., & Smith-Vaughan, H. (2013). Quantitative PCR confirms culture as the gold standard for detection of lower airway infection by nontypeable *Haemophilus influenzae* in Australian Indigenous children with bronchiectasis. *Journal of Microbiological Methods*, 92(3), 270-272. <https://doi.org/10.1016/j.mimet.2012.12.013>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Quantitative PCR confirms culture as the gold standard for detection of lower airway infection by nontypeable *Haemophilus influenzae* in Australian Indigenous children with bronchiectasis {REVISED}

Kim M. Hare^{a,*}, Robyn L. Marsh^a, Michael J. Binks^a, Keith Grimwood^{b,c}, Susan J. Pizzutto^a, Amanda J. Leach^a, Anne B. Chang^{a,b,d,e}, Heidi C. Smith-Vaughan^a

^a *Child Health Division, Menzies School of Health Research, Charles Darwin University, PO Box 41096, Casuarina, Darwin, Northern Territory 0811, Australia*

^b *Queensland Children's Medical Research Institute, The University of Queensland, Royal Children's Hospital, Herston Road, Brisbane, Queensland 4029, Australia*

^c *Department of Infectious Diseases, Royal Children's Hospital, Herston Road, Brisbane, Queensland 4029, Australia*

^d *Royal Darwin Hospital, Rocklands Drive, Tiwi, Northern Territory 0810, Australia*

^e *Department of Respiratory Medicine, Royal Children's Hospital, Herston Road, Brisbane, Queensland 4029, Australia*

*** Corresponding author:**

Email kim.hare@menzies.edu.au

PO Box 41096, Casuarina, Darwin, Northern Territory 0811, Australia

Tel: +61 8 89228196

Fax: +61 8 89275187

Abstract

Correlation was observed between quantitative PCR and semi-quantitative culture for definition of *Haemophilus influenzae* infection in bronchoalveolar lavage specimens from 81 children with bronchiectasis. However, qPCR data correlated less well with airway neutrophilia, and supports continued use of culture as the gold standard for defining *H. influenzae* lower airway infection.

Keywords

Airway cellularity; Bronchiectasis; Estimates of infection; *Haemophilus haemolyticus*; Lower airway infection; Nontypeable *Haemophilus influenzae*;

Nontypeable *Haemophilus influenzae* (NTHi) is frequently cultured from the lower airways of children with bronchiectasis (Hare et al., 2010). However, culture-based phenotypic identification methods cannot reliably differentiate NTHi from nonhemolytic strains of the primarily commensal *Haemophilus haemolyticus*.

Previously, we used a probe-based PCR targeting the protein D gene (*hpd#3*) (Binks et al., 2012; Wang et al., 2011) to differentiate *H. influenzae* and *H. haemolyticus* isolated from bronchoalveolar lavage (BAL) specimens from Australian Indigenous children with bronchiectasis (Hare et al., 2012a). Of 84 BAL specimens tested, 30 (36%) were culture-positive (defined by semi-quantitative growth scores) for NTHi infection. At least one of four isolates collected from these positive specimens was confirmed as *H. influenzae* by *hpd#3* PCR. However, 27% of culture-positive BAL specimens (8/30) had at least one phenotypic NTHi isolate test negative by *hpd#3* PCR, indicating *H. haemolyticus* (Hare et al., 2012a).

Co-detection of these species raised doubts about previous culture-based estimates of infection attributed to *H. influenzae*. Semi-quantitative counts are made on primary plates where *H. influenzae* and *H. haemolyticus* colonies cannot be differentiated, potentially leading to overestimation of *H. influenzae* density. We thus expanded on our previous work by using the *hpd#3* PCR quantitatively (*hpd#3* qPCR) to further define the presence of NTHi in paediatric bronchiectasis. For 81 of the original 84 BAL specimens we compared: (i) the semi-quantitative NTHi culture-based definition with *hpd#3* qPCR estimation of *H. influenzae* infection; and (ii) both measures of bacterial density with total and differential cell counts to gauge the airway inflammatory response to infection.

BAL specimens were collected from children (63% male) aged 5-to-155 months undergoing bronchoscopy between July 2007 and August 2010. Fifty-three children had received antibiotics in the 2-weeks preceding bronchoscopy. BAL specimens were collected, stored and processed (10 μ L aliquots plated on selective media) as described previously (Hare et al., 2010). Growth on primary plates was quantified as: 0=no colonies, 1=1-20 colonies, 2=21-50 colonies, 3=51-100 colonies, 4=>100 colonies within primary inoculum only, 5=growth extending into first streak zone, 6=growth in second streak zone, 7=growth in third streak zone. Phenotypic NTHi was identified by colonial morphology, X and V growth factor requirement, and failure to react with *H. influenzae* types a-f antisera (Phadebact®, Bactus AB, Sweden). Lower airway infection was defined by semi-quantitative growth scores ≥ 4 , which correlate with $>10^4$ colony-forming units (CFU)/mL BAL fluid (Hare et al., 2010).

DNA was extracted from 100 μ L of BAL specimen using QIAamp columns (QIAGEN, Australia) with enzymatic pre-treatment as detailed elsewhere (Smith-Vaughan et al., 2006). The qPCR assay included 5 DNA standards (*H. influenzae* reference isolate ATCC 19418) ranging from 10 to 100,000 genome copies per reaction and was performed as described previously (Binks et al., 2012) using published *hpd#3* primer and probe concentrations (Wang et al., 2011) with TaqMan® Universal PCR Master Mix reagents (Applied Biosystems). All samples, standards and ‘no template’ controls were run in duplicate and required to differ by less than 0.5 cycles. Standard curve correlation coefficients were >0.99 . The qPCR efficiency was ≥ 0.86 . The lower limit of detection was 10 cells (genome copies)/reaction. The lower limit of quantification (LOQ) was 100 cells/reaction (2.5×10^4 cells/mL). Raw data were analysed using Rotor-Gene 6000 Series Software 1.7.

Total and differential cell counts (reflecting lower airway inflammation) were determined using unfiltered BAL specimens as described previously (Pizzutto et al., 2012). Statistical analyses were performed using nonparametric tests in Stata version 10. The Human Research Ethics Committee of the Northern Territory Department of Health and Menzies School of Health Research approved the study (HREC 07/63).

Twenty-nine of the 81 BAL specimens (36%) had phenotypic NTHi growth scores ≥ 4 indicating lower airway infection. Typeable *H. influenzae* strains were found in three specimens; all had concurrent phenotypic NTHi infection. Using qPCR, 40/81 (49%) specimens had *H. influenzae* infection defined as $\geq 10^5$ cells/mL (Abdeldaim et al., 2010). Semi-quantitative growth scores correlated well with \log_{10} qPCR bacterial load estimates (Figure); Spearman's rank correlation coefficient [r_s] was 0.66 ($P < 0.001$) for the 47 specimens with quantifiable PCR results. Of the 34 BAL specimens either *H. influenzae* PCR-negative or positive below the LOQ ($< 2.5 \times 10^4$ cells/mL), 33 had NTHi growth scores of 0 or 1 indicating negative or very low colony counts (< 20 colonies/10 μ L, $< 2 \times 10^3$ CFU/mL).

Twenty-seven of the 29 specimens with NTHi growth scores ≥ 4 had lower airway infection confirmed by qPCR ($\geq 10^5$ cells/mL), including all specimens with concurrent *H. influenzae* and *H. haemolyticus* detection. Thirty-nine specimens were negative for *H. influenzae* infection by both culture (growth score < 4) and qPCR ($< 10^5$ cells/mL). The remaining 15 specimens had discordant culture and qPCR results, including 13 qPCR-positive, culture-negative (growth score < 4) specimens

(Table 1 and Figure). Six of these 13 children had received antibiotics within the previous 2-weeks.

Airway cellularity data were available for 71 (69 with neutrophil data) of the 81 children with BAL culture and qPCR results (Table 2). Total cell counts, neutrophil numbers and neutrophil percentages correlated with both semi-quantitative growth scores (r_s 0.45, 0.48 and 0.41 respectively, all $P < 0.001$) and \log_{10} qPCR values (r_s 0.48 [$P = 0.002$], 0.38 [$P = 0.020$] and 0.31 [$P = 0.057$] respectively). Not unexpectedly (Kapur et al., 2012), airway cellularity measures were significantly higher in children with culture and qPCR-defined *H. influenzae* infection than children without infection by these measures (Table 2). In contrast, airway cellularity was similar between those with *H. influenzae* infection defined solely by qPCR and those infection-negative by both qPCR and culture; inflammatory indices were low in both groups (Table 2).

Several factors may lead to PCR-positive, culture-negative results, such as the presence of non-viable bacteria and the fact that CFU/mL does not invariably correspond to cells/mL since several bacteria may aggregate and generate one colony (Abdeldaim et al., 2010). Additionally, culture sensitivity may be negatively influenced by commensal bacteria numbers and/or antibiotic treatment before sampling (Kais et al., 2006). In our study, most specimens with culture and qPCR-defined infection had commensal bacteria (mainly alpha-hemolytic streptococci) growth scores ≥ 4 , suggesting the presence of commensal organisms did not impede NTHi recovery. Consistent with our earlier report (Hare et al., 2012b), we also found that recent antibiotics were not associated with reduced culture evidence of NTHi lower airway infection. While qPCR had greater sensitivity than culture, the

cellularity data raise questions over the clinical relevance of PCR-positive but culture-negative results.

In conclusion, all BAL specimens with concurrent NTHi and *H. haemolyticus* detection had culture-defined NTHi infection confirmed by *H. influenzae* qPCR. qPCR is consistently reported as more sensitive than culture and is an attractive method for *H. influenzae* quantification. However, the lack of airway inflammation in BAL specimens with *H. influenzae* infection defined by qPCR but not culture suggests that reliance on qPCR data alone may lead to overestimation of lower airway infection rates. While PCR remains useful for determining *H. influenzae* presence and density, our data support continued use of culture as the gold standard for diagnosing NTHi lower airway infection.

ACKNOWLEDGEMENTS

We thank the children and their carers for participating in the study, Menzies staff (Gabrielle McCallum and Kobi Schutz) and Royal Darwin Hospital staff for assistance with specimen collection, and Menzies laboratory staff (Vanya Hampton, Joanna Bugg, Estelle Carter and Peter Christensen) for assistance with laboratory processing. This study was supported by NHMRC project grant 545223.

ABC is supported by NHMRC Practitioner Fellowship 545216. AJL is supported by NHMRC Elizabeth Blackburn Research Fellowship 1020561. HSV is supported by NHMRC Career Development Fellowship 1024175. RM is supported by NHMRC Frank Fenner Early Career Fellowship 1034703. KMH is supported by NHMRC Gustav Nossal Postgraduate Scholarship 1038072. MJB and SJP are supported by NHMRC Postgraduate Scholarships 1017225 and 1038415.

REFERENCES

- Abdeldaim, G.M.K., Stralin, K., Korsgaard, J., Blomberg, J., Welinder-Olsson, C., Herrmann, B., 2010. Multiplex quantitative PCR for detection of lower respiratory tract infection and meningitis caused by *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Neisseria meningitidis*. *BMC Microbiol.* 10, 310.
- Binks, M.J., Temple, B., Kirkham, L.A., Wiertsema, S.P., Dunne, E.M., Richmond, P.C., Marsh, R., Leach, A.J., Smith-Vaughan, H., 2012. Molecular surveillance of true nontypeable *Haemophilus influenzae*: An evaluation of PCR screening assays. *PLoS One* 7, e34083.
- Hare, K.M., Grimwood, K., Leach, A.J., Smith-Vaughan, H., Torzillo, P.J., Morris, P.S., Chang, A.B., 2010. Respiratory bacterial pathogens in the nasopharynx and lower airways of Australian Indigenous children with bronchiectasis. *J. Pediatr.* 157, 1001-1005.
- Hare, K.M., Binks, M.J., Grimwood, K., Chang, A.B., Leach, A.J., Smith-Vaughan, H., 2012a. Culture and PCR detection of *Haemophilus influenzae* and *Haemophilus haemolyticus* in Australian Indigenous children with bronchiectasis. *J. Clin. Microbiol.* 50, 2444-2445.
- Hare, K.M., Leach, A.J., Morris, P.S., Smith-Vaughan, H., Torzillo, P., Bauert, P., Cheng, A.C., McDonald, M.I., Brown, N., Chang, A.B., Grimwood, K., 2012b. Impact of recent antibiotics upon nasopharyngeal carriage and lower airway infection in Indigenous Australian children with non-CF bronchiectasis. *Int. J. Antimicrobial Agents* 40, 365-369.
- Kais, M., Spindler, C., Kalin, M., Ortqvist, A., Giske, C.G., 2006. Quantitative detection of *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Moraxella*

catarrhalis in lower respiratory tract samples by real-time PCR. *Diagn. Microbiol. Infect. Dis.* 55, 169-178.

Kapur, N., Grimwood, K., Masters, B., Morris, P.S., Chang, A.B., 2012. Lower airway microbiology and cellularity in children with newly diagnosed non-CF bronchiectasis. *Pediatr. Pulmonol.* 47, 300-307.

Pizzutto, S.J., Grimwood, K., Bauert, P., Schultz, K.L., Yerkovich, S.T., Upham, J.T., Chang, A.B., 2012. Bronchoscopy contributes to the clinical management of Indigenous children newly diagnosed with bronchiectasis. *Pediatr. Pulmonol.* Mar 19 [Epub ahead of print]

Smith-Vaughan, H., Byun, R., Nadkarni, M., Jacques, N.A., Hunter, N., Halpin, S., Morris, P.S., Leach, A.J., 2006. Measuring nasal bacterial load and its association with otitis media. *BMC Ear Nose Throat Disord.* 6, 10.


Wang, X., Mair, R., Hatcher, C., Theodore, M.J., Edmond, K., Wu, H.M., Harcourt, B.H., Carvalho, M.D., Pimenta, F., Nymadawa, P., Altantsetseg, D., Kirsch, M., Satola, S.W., Cohn, A., Messonnier, N.E., Mayer, L.W., 2011. Detection of bacterial pathogens in Mongolia meningitis surveillance with a new real-time PCR assay to detect *Haemophilus influenzae*. *Int. J. Med. Microbiol.* 301, 303-309.

Table 1: *Haemophilus influenzae* lower airway infection in Australian Indigenous children defined by semi-quantitative culture and quantitative PCR

Growth score ≥ 4	qPCR $\geq 10^5$ cells/mL		Total
	Negative	Positive	
Negative	39	13	52
Positive	2	27	29
Total	41	40	81

Using culture as the gold standard, qPCR sensitivity is 93% and specificity 75%.

Figure: *Haemophilus influenzae* lower airway infection in Australian Indigenous children defined by semi-quantitative culture and quantitative PCR



Scatter plot with linear trend line showing culture-based semi-quantitative growth scores of phenotypic nontypeable *Haemophilus influenzae* (NTHi) and log₁₀ *Haemophilus influenzae* hpd#3 qPCR results from 47 bronchoalveolar lavage (BAL) specimens with a quantifiable specific PCR result. The oval encompasses specimens with lower airway infection as determined by semi-quantitative culture (score ≥ 4) and qPCR ($\geq 10^5$ cells/mL).

Table 2: Airway cellularity data for Australian Indigenous children with and without *Haemophilus influenzae* lower airway infection^a

Culture-defined NTHi infection	Negative (growth score <4)	Positive (growth score ≥4)	P-value^b
Median (IQR) TCC (x 10⁶ cells/L)	N=49 350 (240, 570)	N=22 810 (460, 1210)	P < 0.001
Median (IQR) neutrophils (x 10⁶ cells/L)	N=48 30 (10, 120)	N=21 250 (100, 740)	P = 0.002
Median (IQR) neutrophil percentage	N=47 11 (6, 31)	N=22 35 (9, 62)	P = 0.022
qPCR-defined Hi infection	Negative (<10⁵ cells/mL)	Positive (≥10⁵ cells/mL)	P-value^b
Median (IQR) TCC (x 10⁶ cells/L)	N=38 340 (200, 580)	N=33 650 (370, 1020)	P = 0.003
Median (IQR) neutrophils (x 10⁶ cells/L)	N=37 30 (10, 140)	N=32 120 (30, 600)	P = 0.005
Median (IQR) neutrophil percentage	N=36 10 (3, 32)	N=33 21 (11, 59)	P = 0.015

qPCR-defined Hi infection for NTHi culture negative BALs	Culture negative (growth score <4)	Culture negative (growth score <4)	P-value^b
	qPCR negative (<10⁵ cells/mL)	qPCR positive (≥10⁵ cells/mL)	
Median (IQR) TCC (x 10⁶ cells/L)	N=36 340 (190, 560)	N=13 370 (290, 570)	P = 0.497
Median (IQR) neutrophils (x 10⁶ cells/L)	N=35 30 (10, 140)	N=13 50 (20, 100)	P = 0.190
Median (IQR) neutrophil percentage	N=34 10 (2, 31)	N=13 13 (11, 28)	P = 0.171

BAL, bronchoalveolar lavage; Hi, *Haemophilus influenzae*; IQR, interquartile range;

NTHi, nontypeable Hi; qPCR, quantitative PCR; TCC, total cell count

^a Exclusion of 21 BAL samples with growth scores ≥4 for respiratory pathogens other than Hi did not alter these findings

^b Two-sample Wilcoxon rank-sum (Mann-Whitney) test