



Charles Darwin University

Technical note

Dynamic INtegrated Gap-filling and partitioning for OzFlux (DINGO)

Beringer, Jason; McHugh, Ian; Hutley, Lindsay B.; Isaac, Peter; Kljun, Natascha

Published in:
Biogeosciences

DOI:
[10.5194/bg-14-1457-2017](https://doi.org/10.5194/bg-14-1457-2017)

Published: 23/03/2017

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Beringer, J., McHugh, I., Hutley, L. B., Isaac, P., & Kljun, N. (2017). Technical note: Dynamic INtegrated Gap-filling and partitioning for OzFlux (DINGO). *Biogeosciences*, 14(6), 1457-1460. <https://doi.org/10.5194/bg-14-1457-2017>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Technical note: Dynamic INtegrated Gap-filling and partitioning for OzFlux (DINGO)

Jason Beringer¹, Ian McHugh², Lindsay B. Hutley³, Peter Isaac⁴, and Natascha Kljun⁵

¹School of Earth and Environment (SEE), The University of Western Australia, Crawley WA, 6009, Australia

²School of Earth, Atmosphere and Environment, Monash University, Clayton, 3800, Australia

³School of Environment, Research Institute for the Environment and Livelihoods, Charles Darwin University, NT 0909, Australia

⁴School of Earth, Atmosphere and Environment, Monash University, Clayton, 3800, Australia

⁵Department of Geography, Swansea University, Singleton Park, Swansea, Wales SA2 8PP, UK

Correspondence to: Jason Beringer (jason.beringer@uwa.edu.au)

Received: 5 May 2016 – Discussion started: 10 May 2016

Revised: 7 February 2017 – Accepted: 7 February 2017 – Published: 23 March 2017

Abstract. Standardised, quality-controlled and robust data from flux networks underpin the understanding of ecosystem processes and tools necessary to support the management of natural resources, including water, carbon and nutrients for environmental and production benefits. The Australian regional flux network (OzFlux) currently has 23 active sites and aims to provide a continental-scale national research facility to monitor and assess Australia’s terrestrial biosphere and climate for improved predictions. Given the need for standardised and effective data processing of flux data, we have developed a software suite, called the Dynamic INtegrated Gap-filling and partitioning for OzFlux (DINGO), that enables gap-filling and partitioning of the primary fluxes into ecosystem respiration (Fre) and gross primary productivity (GPP) and subsequently provides diagnostics and results. We outline the processing pathways and methodologies that are applied in DINGO (v13) to OzFlux data, including (1) gap-filling of meteorological and other drivers; (2) gap-filling of fluxes using artificial neural networks; (3) the u^* threshold determination; (4) partitioning into ecosystem respiration and gross primary productivity; (5) random, model and u^* uncertainties; and (6) diagnostic, footprint calculation, summary and results outputs. DINGO was developed for Australian data, but the framework is applicable to any flux data or regional network. Quality data from robust systems like DINGO ensure the utility and uptake of the flux data and facilitates synergies between flux, remote sensing and modelling.

1 Introduction

OzFlux is the regional Australian and New Zealand flux tower network that aims to provide a continental-scale national research facility to monitor and assess Australia’s terrestrial biosphere and climate for improved predictions (Beringer et al., 2016). High-quality and reliable data are a crucial foundation in achieving the objectives of the OzFlux network (Beringer et al., 2016) and underpin the process understanding needed to (1) support sound management of natural resources including water, carbon and nutrient resources for environmental and production benefits; (2) monitor, assess, predict and respond to climate change and variability; (3) improve weather and environmental information and prediction; (4) support disaster management and early warning systems needed to meet Australia’s priorities in national security; and (5) ensure that Earth system models used to underpin Australia’s policies and commitments to international treaties adequately represent Australian terrestrial ecosystem processes (Beringer et al., 2016).

Beringer et al. (2016) provide an overview of the evolution, design and current status of OzFlux as well as a brief summary of the instrumentation and data collection that forms the backbone of the network. A detailed description of the quality control and post-processing of the eddy covariance data using OzFluxQC and the data pathway to curation is given by Isaac et al. (2016). In summary, from Beringer et al. (2016), most sites have data loggers that provide the average (usually over 30 min) covariances that are processed

through 6 levels using the OzFluxQC standard software processing scripts. Levels 1, 2 and 3 are as follows: L1 is the raw data as received from the flux tower, L2 is the quality-controlled data, and L3 is the post-processed, corrected, but not gap-filled data. Quality control measures by OzFluxQC are applied at L2 and comprise checks for plausible value ranges, spike detection and removal, manual exclusion of date and time ranges and diagnostic checks for all quantities used in the flux correction calculations. The quality checks make use of the diagnostic information from the sonic anemometer and the infrared gas analyser. For sites calculating fluxes from the averaged covariances, post-processing includes 2-D coordinate rotation, low- and high-pass frequency correction, conversion of virtual heat flux to sensible heat flux and application of the Webb–Pearman–Leuning (WPL) correction to the latent heat and CO₂ fluxes (see Burba, 2013 for a general description of the data processing pathways). Steps performed at L3 include the correction of the ground heat flux for storage in the layer above the heat flux plates (Mayocchi and Bristow, 1995) and correction of the CO₂ flux data for storage in the canopy (where available). OzFlux data are available at <http://data.ozflux.org.au/>. OzFlux sites submit their data to FLUXNET at L3.

Given the international need by the community for standardised data processing to enable effective comparison across biomes and to understand inter-annual variability (Papale et al., 2006), we have developed a software tool to address this need. In this paper, we describe the development and testing of the Dynamic INtegrated Gap-filling and partitioning for the Ozflux (DINGO) system that utilises the L3 data from OzFluxQC to gap-fill and partition the fluxes in ecosystem respiration (Fre) and gross primary productivity (GPP) and subsequently provides diagnostics and results. This paper is not intended to be a thorough review of the data processing, but the application of standard techniques in DINGO for the flux community. DINGO is a research version for OzFlux data, whereas the OzFluxQC system (which has many similar features to DINGO) is considered an operational version. Quality data from robust systems like DINGO ensure the utility and uptake of the flux data and facilitates synergies between flux, remote sensing, modelling and canopy physiological studies. We conclude by looking ahead at the future direction of the DINGO system.

2 Approach

The overall approach used in DINGO is to take the L3 OzFluxQC data – which has gaps from data processing (data excluded due to values out of range, spike detection or manual exclusion of date and time ranges) or from site issues (instrument or power failure, herbivores, fire, eagles nests, cows, lightning, PI on sabbatical, etc.) – and gap-fill and partition the data using a variety of data sources (Fig. S1). DINGO is programmed in python 2.7 and is currently at ver-

sion 13 and publically available on GitHub (<https://github.com/jberinge/DINGO13>). It is designed to work with OzFlux data produced in NetCDF format by the OzFluxQC (Isaac et al., 2016) and draws on Australian automatic weather station (AWS) data, but it could be adapted for other data sources across other flux sites. The primary interface for the user is through a text-based control file that has information on site characteristics (name, latitude and longitude, the frequency of the flux measurements, i.e. 30 or 60 min, and elevation, file paths (to the OzFluxQC NetCDF files and other ancillary data inputs), data processing options, and data plotting and output formats. In general, prior to the processing steps below, any gaps in fluxes or meteorological quantities of less than 2 h are filled by DINGO using linear interpolation. The pathway for processing is shown in Fig. S1, and each step is outlined in the Supplement.

3 Conclusions

The OzFlux network has been highly successful in generating standardised measurements and protocols that provide robust primary data. Only via transparent, advanced and consistent QA/QC will we ensure compatibility within the OzFlux network (Beringer et al., 2016) and with international databases (FLUXNET) (Papale et al., 2006), ensuring uptake by the broader scientific community. Through robust software systems such as OzFluxQC (Isaac et al., 2016) and DINGO, we are able to ensure timely and quality gap-filling and partitioning of fluxes that in turn enable a significant uptake of the eddy covariance data for application to a range of research questions as exemplified in Beringer et al. (2016). This includes integration of the eddy covariance and remote sensing datasets for the validation of satellite products (e.g. GPP and ET, Kanniah et al., 2009; Restrepo-Coupe et al., 2016) and to aid the parameterisation of models that rely on remotely sensed data (e.g. GPP, ET, canopy conductance, and light use efficiency (LUE), Barraza et al., 2014, 2015; Glenn et al., 2011; Goerner et al., 2011). In addition, OzFlux data have been instrumental in constraining a continent-wide assessment of terrestrial carbon and water cycles (Haverd et al., 2013) and featured in the development of new models (Haverd et al., 2007, 2009). There is utility of the data to support carbon accounting activities (Hutley et al., 2005), as demonstrated in research focussed on the conversion of savanna to pasture (Bristow et al., 2016). Ultimately, flux data are required to address the key ecosystem science questions of OzFlux (Beringer et al., 2016) that are focused on the improved understanding of the responses of carbon and water cycles of Australian ecosystems to current climate and disturbance regimes as well as on impacts of projected future changes to precipitation, temperature and atmospheric CO₂ concentration. Key questions include the following: (1) what are the key drivers of ecosystem productivity (carbon sinks) and greenhouse gas emissions; (2) how resilient is the Aus-

tralian ecosystem productivity to an increasingly variable and changing climate; and, (3) what is the current water budget of the dominant Australian ecosystems and how will it change in the future?

Data availability. The data used to create the example figures in the manuscripts were from two sites in the OzFlux network. Site data for Whroo can be found in Beringer (2013) “Whroo OzFlux tower site OzFlux: Australian and New Zealand Flux Research and Monitoring” <http://hdl.handle.net/102.100.100/14232>, whilst the Calperum data can be found in Calperum Tech (2013) “Calperum Chowilla OzFlux tower site OzFlux: Australian and New Zealand Flux Research and Monitoring” <http://hdl.handle.net/102.100.100/14236>.

The Supplement related to this article is available online at doi:10.5194/bg-14-1457-2017-supplement.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work utilised data collected by grants funded by the Australian Research Council DP130101566. Beringer is funded under an ARC Future Fellowship (FT110100602).

Edited by: D. Papale

Reviewed by: three anonymous referees

References

- Barraza, V., Grings, F., Ferrazzoli, P., Huete, A., Restrepo-Coupe, N., Beringer, J., Van Gorsel, E., and Karszenbaum, H.: Behavior of multitemporal and multisensor passive microwave indices in Southern Hemisphere ecosystems, *J. Geophys. Res.–Biogeosci.*, 119, 2231–2244, doi:10.1002/2014JG002626, 2014.
- Barraza, V., Restrepo-Coupe, N., Huete, A., Grings, F., and Van Gorsel, E.: Passive microwave and optical index approaches for estimating surface conductance and evapotranspiration in forest ecosystems, *Agric. For. Meteorol.*, 213, 126–137, doi:10.1016/j.agrformet.2015.06.020, 2015.
- Beringer, J.: Whroo OzFlux tower site, available at: <http://hdl.handle.net/102.100.100/14232> (last access: March 2017), 2013.
- Beringer, J., Hutley, L. B., McHugh, I., Arndt, S. K., Campbell, D., Cleugh, H. A., Cleverly, J., Resco de Dios, V., Eamus, D., Evans, B., Ewenz, C., Grace, P., Griebel, A., Haverd, V., Hinko-Najera, N., Huete, A., Isaac, P., Kanniah, K., Leuning, R., Liddell, M. J., Macfarlane, C., Meyer, W., Moore, C., Pendall, E., Phillips, A., Phillips, R. L., Prober, S. M., Restrepo-Coupe, N., Rutledge, S., Schroder, I., Silberstein, R., Southall, P., Yee, M. S., Tapper, N. J., van Gorsel, E., Vote, C., Walker, J., and Wardlaw, T.: An introduction to the Australian and New Zealand flux tower network – OzFlux, *Biogeosciences*, 13, 5895–5916, doi:10.5194/bg-13-5895-2016, 2016.
- Bristow, M., Hutley, L. B., Beringer, J., Livesley, S. J., Edwards, A. C., and Arndt, S. K.: Quantifying the relative importance of greenhouse gas emissions from current and future savanna land use change across northern Australia, *Biogeosciences*, 13, 6285–6303, doi:10.5194/bg-13-6285-2016, 2016.
- Burba, G.: Eddy Covariance Method for Scientific, Industrial, Agricultural, and Regulatory Applications: A Field Book on Measuring Ecosystem Gas Exchange and Areal Emission Rates, LI-COR Biosciences, Lincoln, NE, USA, 2013.
- Calperum Tech: Calperum Chowilla OzFlux tower site, available at: <http://hdl.handle.net/102.100.100/14236> (last access: March 2017), 2013.
- Glenn, E. P., Doody, T. M., Guerschman, J. P., Huete, A. R., King, E. A., McVicar, T. R., Van Dijk, A. I. J. M., Van Niel, T. G., Yebra, M., and Zhang, Y.: Actual evapotranspiration estimation by ground and remote sensing methods: the Australian experience, *Hydrol. Process.*, 25, 4103–4116, doi:10.1002/hyp.8391, 2011.
- Goerner, A., Reichstein, M., Tomelleri, E., Hanan, N., Rambal, S., Papale, D., Dragoni, D., and Schimmlus, C.: Remote sensing of ecosystem light use efficiency with MODIS-based PRI, *Biogeosciences*, 8, 189–202, doi:10.5194/bg-8-189-2011, 2011.
- Haverd, V., Cuntz, M., Leuning, R., and Keith, H.: Air and biomass heat storage fluxes in a forest canopy: Calculation within a soil vegetation atmosphere transfer model, *Agric. For. Meteorol.*, 147, 125–139, 2007.
- Haverd, V., Leuning, R., Griffith, D., van Gorsel, E., and Cuntz, M.: The Turbulent Lagrangian Time Scale in Forest Canopies Constrained by Fluxes, Concentrations and Source Distributions, *Boundary-Layer Meteorol.*, 130, 209–228, doi:10.1007/s10546-008-9344-4, 2009.
- Haverd, V., Raupach, M. R., Briggs, P. R., Canadell, J. G., Isaac, P., Pickett-Heaps, C., Roxburgh, S. H., van Gorsel, E., Viscarra Rossel, R. A., and Wang, Z.: Multiple observation types reduce uncertainty in Australia’s terrestrial carbon and water cycles, *Biogeosciences*, 10, 2011–2040, doi:10.5194/bg-10-2011-2013, 2013.
- Hutley, L. B., Leuning, R., Beringer, J., and Cleugh, H. A.: The utility of the eddy covariance techniques as a tool in carbon accounting: tropical savanna as a case study, *Aust. J. Bot.*, 53, 663–675, doi:10.1071/BT04147, 2005.
- Isaac, P. R., Cleverly, J., Beringer, J., and McHugh, I.: The OzFlux network data path: from collection to curation, to be submitted to *Biogeosciences Discuss.*, 2016.
- Kanniah, K. D., Beringer, J., Hutley, L. B., Tapper, N. J., and Zhu, X.: Evaluation of Collections 4 and 5 of the MODIS Gross Primary Productivity product and algorithm improvement at a tropical savanna site in northern Australia, *Remote Sens. Environ.*, 113, 1808–1822, doi:10.1016/j.rse.2009.04.013, 2009.
- Mayocchi, C. L. and Bristow, K. L.: Soil surface heat flux: some general questions and comments on measurements, *Agric. For. Meteorol.*, 75, 43–50, 1995.
- Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., Longdoz, B., Rambal, S., Valentini, R., Vesala, T., and Yakir, D.: Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation, *Biogeosciences*, 3, 571–583, doi:10.5194/bg-3-571-2006, 2006.

Restrepo-Coupe, N., Huete, A., Davies, K., Cleverly, J., Beringer, J., Eamus, D., van Gorsel, E., Hutley, L. B., and Meyer, W. S.: MODIS vegetation products as proxies of photosynthetic potential along a gradient of meteorologically and biologically driven ecosystem productivity, *Biogeosciences*, 13, 5587–5608, doi:10.5194/bg-13-5587-2016, 2016.

Schmidt, A., Hanson, C., Chan, W. S., and Law, B. E.: Empirical assessment of uncertainties of meteorological parameters and turbulent fluxes in the AmeriFlux network, *J. Geophys. Res.*, 117, G04014, doi:10.1029/2012JG002100, 2012.