



Charles Darwin University

Compilation and traits of Australian bird species killed by cats

Woinarski, J. C.Z.; Woolley, L. A.; Garnett, S. T.; Legge, S. M.; Murphy, B. P.; Lawes, M. J.; Comer, S.; Dickman, C. R.; Doherty, T. S.; Edwards, G.; Nankivill, A.; Palmer, R.; Paton, D.

Published in:
Biological Conservation

DOI:
[10.1016/j.biocon.2017.09.017](https://doi.org/10.1016/j.biocon.2017.09.017)

Published: 01/12/2017

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

Woinarski, J. C. Z., Woolley, L. A., Garnett, S. T., Legge, S. M., Murphy, B. P., Lawes, M. J., Comer, S., Dickman, C. R., Doherty, T. S., Edwards, G., Nankivill, A., Palmer, R., & Paton, D. (2017). Compilation and traits of Australian bird species killed by cats. *Biological Conservation*, 216, 1-9.
<https://doi.org/10.1016/j.biocon.2017.09.017>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

Compilation and traits of Australian bird species killed by cats

J.C.Z. Woinarski^{a}, L. A. Woolley^a, S.T. Garnett^a, S.M. Legge^b, B.P. Murphy^a, M.J. Lawes^c, S. Comer^d, C.R. Dickman^e, T.S. Doherty^f, G. Edwards^g, A. Nankivill^h, R. Palmerⁱ, D. Paton^j*

^a NESP Threatened Species Recovery Hub, Research Institute for the Environment and Livelihoods, Charles Darwin University, Casuarina, Northern Territory 0909, Australia.

^b NESP Threatened Species Recovery Hub, Centre for Biodiversity and Conservation Research, University of Queensland, St Lucia, Qld 4072, Australia

^c School of Life Sciences, University of KwaZulu-Natal, Scottsville, 3209, South Africa

^d Department of Biodiversity, Conservation and Attractions, Albany, WA 6330, Australia

^e NESP Threatened Species Recovery Hub, Desert Ecology Research Group, School of Life and Environmental Sciences A08, University of Sydney, NSW 2006, Australia

^f Deakin University, School of Life and Environmental Sciences, Centre for Integrative Ecology (Burwood Campus), Geelong, Australia

^g Northern Territory Department of Land Resource Management, PO Box 1120, Alice Springs, NT 0871, Australia ^h Nature Foundation SA Inc, PO Box 448, Hindmarsh SA 5007

ⁱ Department of Biodiversity, Conservation and Attractions, Locked Bag 104, Bentley Delivery Centre, WA 6983, Australia

^j School of Biological Sciences, University of Adelaide, SA 5005, Australia

*Corresponding author. email: john.woinarski@cdu.edu.au; phone (+61)3 975 44094; 0455961000; postal address: 38 Colby Drive, Belgrave Heights, Victoria, 3160, AUSTRALIA

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70

Abstract. House cats *Felis catus* have contributed to the extinction of many bird species on islands, but their impact on continental bird faunas is less well resolved. Here, we compile and analyse a comprehensive record of all bird species known to be killed by feral cats at a continental scale. From published studies and unpublished data, we document predation by feral and pet cats on 357 bird species in Australia, including 338 Australian (non-vagrant) native bird species (=45.6% of the 741 Australian native bird species, excluding vagrants). This tally included 24 species listed as threatened or extinct by the IUCN (40% of the 58 non-vagrant Australian species listed as threatened), and 71 of the 117 bird species (61%) listed as threatened under Australian legislation (or species with one or more subspecies so listed). These tallies are substantially larger than reported in previous reviews. We provide the first continental-scale attempt to model bird species' traits that are associated with likelihood of being killed by cats, and use such modelling to attempt to redress some inevitable biases in compilation of predation records on birds. We conclude that the likelihood of being killed by a cat was highest for bird species that are restricted to islands, are of intermediate body mass (ca. 60-300 g), and nest and forage on the ground, and least likely for bird species occurring mostly in rainforests and wetlands. We also identify a set of bird species most likely to be threatened by cat-predation and hence most likely to benefit from enhanced management of cats. This study does not specifically evaluate the impact of cats on bird populations or on the conservation of Australian birds, but our results suggest that such impact may be much more pervasive than previously documented.

Running head: Australian bird species killed by cats

Additional key words: diet, invasive predator, modelling, threatened species

71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115

Introduction

Cats *Felis catus* are versatile predators that largely employ an ‘ambush’ hunting strategy (Bradshaw 1992; Turner and Meister 1988) to capture and kill a very wide range of animal species from small invertebrates to vertebrates up to at least 4 kg (Bonnaud et al. 2011; Fancourt 2015). Predation by introduced cats has been a major cause of extinction for many species, with such impact particularly pronounced for island-endemic vertebrates (Blackburn et al. 2004; Blackburn et al. 2005; Doherty et al. 2016; Medina et al. 2011; Nogales et al. 2013) and for mammals in Australia (Woinarski et al. 2015). In contrast, the impacts of predation by cats on continental bird faunas is less well resolved, although cats are known to kill hundreds of millions to billions of birds annually in continental settings (Blancher 2013; Dauphiné and Cooper 2009; Loss et al. 2013), with such predation shown to be a major source of bird mortality (Loss et al. 2012, 2015).

In a recent paper, Woinarski et al. (in press) concluded that about one million birds are killed in Australia per day by cats. However that study provided no information on the extent to which this toll fell equitably or otherwise across bird species. Here, we complement that previous paper by reporting on the Australian bird species known to be killed by cats, and seek to identify bird species, or groupings of species, that are most likely to be subject to cat predation.

First introduced to Australia in the late eighteenth century (Abbott 2008), cats are now almost ubiquitous across the Australian mainland and also occur on many Australian islands (Legge et al. 2017). There have been two notable listings of Australian bird species known to have been preyed upon by cats. An extensive survey of pet-owners in south-eastern Australia reported records of pet cats killing (or capturing) individuals of 186 bird species (Paton 1990; Paton 1991; Paton 1993), although the full list associated with that study has never been formally published. More recently, Doherty et al. (2015) aggregated information from 70 published and unpublished studies, widely spaced across Australia, of the diet of feral cats. That review compiled cat-predation records for 123 bird species, including 113 native species, of which two species were listed by the IUCN as threatened (Malleefowl *Leipoa ocellata* and Southern Rockhopper Penguin *Eudyptes chrysocome*). Another recent but more speculative compilation relating to the possible detrimental effects of feral cats on Australian biodiversity listed 40 threatened Australian bird taxa (including subspecies) that ‘may be affected by predation by feral cats’, although in many of these cases this implication was not based on any definite records of such predation (Department of the Environment 2015).

As recognised by their authors, the lists of bird species reported as preyed upon by cats in these previous compilations have some substantial biases and incompleteness (Table 1). Paton’s set of studies were based on cat-owners’ records in urban and rural areas of south-eastern Australia, and hence bird species that are readily identified by the public were more likely reported by respondents, and bird species that are more common and widespread in this region were likely to have contributed most to the cat-killed tallies. The compilation by Doherty et al. (2015) was more geographically representative, but was also likely to include more common and widespread bird species, and species for which partly-digested prey items are readily identifiable to species. Rare and restricted bird species are less likely to be reported as cat-prey in these data sets, but it is possible that such species have a higher *per capita* rate of being preyed upon by cats, and hence suffer more

116 conservation impact, than those bird species that – because of their abundance or wide distributions
117 – are more likely to be reported in cat dietary studies.

118

119 In the current study, we build on these important preceding compilations through inclusion of
120 records from many additional and more diverse sources in order to provide a continental-scale
121 compilation of bird species for which there are records of individual birds killed by cats, noting also
122 the threatened bird species in this compilation. We then examine, across all Australian bird species,
123 for relationships between records of cat predation and bird species' ecological, morphological and
124 other traits. We then model these relationships to rank species according to their likelihood of being
125 killed by cats, with and without controls for a measure of bird abundance and range. Our modelling
126 at continental scale seeks to diminish the bias due to cat predation being more likely to have been
127 recorded for bird species that are common in areas with higher human population density. This bias
128 may be particularly important to try to redress because a recent continental-scale assessment of
129 predation by feral cats in Australia (Woinarski et al. in press) reported that the modelled rate of
130 predation of birds (i.e. no. individual birds killed km⁻² y⁻¹) by cats was highest in arid and semi-arid
131 areas remote from most human population centres, and hence bird species in those relatively
132 under-studied areas may be most at risk from cat predation.

133

134

135 **Methods**

136

137 Terminology. Note that for convenient shorthand here we use the expression 'bird species killed by
138 cats', or variants. We recognise that it is individuals, rather than species, that are killed; but repeated
139 use of that correct wording is unduly cumbersome.

140

141 *Compilation of cat-predation database*

142 We sought records of birds being killed by cats from many diverse sources. The most notable of
143 these included:

144

- 145 • cat dietary studies (including and extending all sources used in Doherty et al. (2015));
- 146 • a small number of largely anecdotal records compiled in the Handbook of Australian and
147 New Zealand and Antarctic Birds series (Higgins 1999; Higgins and Davies 1996; Higgins and
148 Peter 2002; Higgins et al. 2006; Higgins et al. 2001; Marchant and Higgins 1990, 1993), the
149 compendium of all information then available about Australian birds, in which sources of
150 bird mortality (including "street urchins" and "horseless carriages") are occasionally
151 provided;
- 152 • autecological studies of bird species, where these provided information on causes of
153 mortality (e.g. Smith and Saunders 1986);
- 154 • unpublished records from the Australian Bird and Bat Banding Scheme of reported causes of
155 mortality or injury to banded birds (340 records of cat-killed birds of 124 species);
- 156 • unpublished records from all Australian museums (372 specimen records of cat-killed birds
157 of 110 species); and
- 158 • compilations of injured wildlife reported by veterinarians, where the cause of injury was
159 reported (Dowling et al. 1994).

160

161 A total of 86 published sources (including reports and theses) with records of Australian birds being
162 killed by cats are included in this compilation (Appendix B); augmented by a further 18 unpublished
163 studies that provided information on contents of a total of 1571 cat stomachs or scats (Appendix A).
164 About ten of the published sources are largely secondary, but the distinction between primary and
165 secondary sources was not always clear in the literature. Although some published or unpublished
166 records of birds being killed by cats clearly indicated the subspecies of birds being consumed, most
167 did not, so our compilation is at species level only.

168

169 We include cases of birds known to be injured (but not necessarily killed and consumed) by cats. We
170 include records of cats consuming eggs and nestlings, in the few cases where the bird species was
171 identified. Some of the dietary records may be a result of cats scavenging on dead birds (perhaps
172 especially in the case of larger bird species), but in many cases it is impossible to determine whether
173 items reported in a cat's stomach or scat are a result of predation or scavenging. In general, cats
174 prefer hunting live prey to scavenging, but they are known to consume carrion (Doherty et al. 2015;
175 Molsher et al. 2017).

176

177 Some of the literature we searched incidentally included records of cats killing birds of species that
178 occur in Australia, but for which the reported predation occurred outside Australia. We noted these
179 records (in Appendix B), but we have not attempted to review literature of cat-predation beyond
180 Australia, and we do not include these records in our analyses or tallies. Some sources also noted
181 that cat-predation was inferred, rather than being supported by definitive evidence. Such records
182 are noted in Appendix B as inferred predation, but are also not included in our tallies or modelling.

183

184 This compilation does not differentiate between predation by pet or feral cats because a substantial
185 proportion of the primary sources that we examined did not make this distinction. Furthermore,
186 there is a continuum from, at one extreme, pet cats that are not allowed outside (for which all food
187 is provided by their human owners) to, at the other extreme, feral cats in natural environments
188 remote from humans.

189

190 *Bird species traits*

191 Our listing of Australian bird species was from the recent comprehensive data base of Garnett et al.
192 (2015): these include species occurring on the Australian mainland and islands, including Australia's
193 overseas territories. That source also categorised some of these species as vagrant, and unless
194 otherwise indicated, such species are omitted from analyses here. The threatened status of every
195 bird species as at January 2017 was also included in our database, at both global level (i.e. by the
196 IUCN) and national level (as recognised by Australia's *Environment Protection and Biodiversity
197 Conservation Act, 1999*). Note that the Australian legislation allows listing of subspecies as
198 threatened; in this study, we report killing by cats only at the species level, but if a cat is known to
199 kill one subspecies of a particular bird species, it is reasonable to assume that it is likely to also kill
200 another subspecies of that species.

201

202 For every bird species, we tallied the number of different sources that reported predation by cats.
203 We also condensed this to a binary variable – whether there were or were not confirmed records of
204 cat-predation in Australia for that species in our collated database. We also compiled a set of
205 ecological, morphological and other variables for every Australian bird species (Table 2), with traits

206 included based largely on results from previous studies that have indicated some factors associated
207 with the likelihood of a bird species being preyed upon by cats, including body mass, nest site and
208 foraging substrate (Dickman 1996; Kutt 2012; Lepczyk et al. 2004; Paltridge et al. 1997). Our scoring
209 for these factors was mostly derived from the comprehensive database of traits of Australian birds
210 (Garnett et al. 2015), although some were simplified from that source to provide tractability in the
211 modelling (see Appendix C). We could not readily derive, and hence do not include in modelling,
212 information about some additional traits that may also differentially affect the likelihood of a bird
213 species being preyed upon by cats. For example, scent may be important for some mammalian
214 predators, and some bird species (e.g. Ground Parrots *Pezoporus wallicus*) are considered
215 particularly detectable to mammalian predators because of their strong scent (Mattingley 1918).
216 Likewise, bird species that have conspicuously marked plumage may also be more readily detected
217 by hunting cats; some bird species may be characteristically more wary than others; and some bird
218 species may respond vigorously and pugnaciously to attempted attacks.

219

220 For every bird species, we also included two variables that relate to their abundance, distribution
221 and the extent to which the species has been subject to research. The variables were: (i) the number
222 of observations reported in the two Atlases of Australian Birds (1977 to 1981, and 1998 to 2001)
223 combined. This value will tend to be higher for species that are more widespread and abundant,
224 with substantial distributions overlapping that of major human population centres (i.e. where most
225 observers reside). For idiosyncratic reasons, the Atlas tallies do not include any records from oceanic
226 islands; and (ii) the number of individual birds banded, a measure of targeted research effort, which
227 again is likely to be higher for species that are more widespread and abundant, with substantial
228 distribution overlapping that of major human population centres, but may also be high for some
229 rarer and more restricted species that have happened to have been subject to intensive research
230 programs. Given that there is more information available, including more targeted studies, for
231 species with higher values for these variables, it is likely that species with high values for these
232 variables will be more likely to have documented records of being killed by cats than would
233 otherwise similar bird species that have low values for these variables (i.e. are rarer, more restricted
234 or less studied), even though their *per capita* rate of predation by cats may be comparable. In
235 analyses (below) we seek to redress this bias.

236

237 *Analysis*

238 As one approach to considering the extent to which our compilation of diverse sources redresses
239 potential bias arising from common and widespread bird species being particularly likely to be
240 reported in cat dietary studies, we compared the abundance and distributional extent of the set of
241 bird species recorded as killed by cats in the Doherty et al. (2015) compilation, the set of additional
242 bird species recorded here as killed by cats, and the set of bird species that have not yet been
243 reported to be killed by cats, using Kruskal-Wallis analysis of variance.

244

245 Our principal analysis modelled the presence/absence of cat-predation records for Australian bird
246 species against all possible combinations of bird species' traits using generalized linear models
247 (GLM's) (binomial logistic regression) run in R version 3.3.2 (R Core Team 2016). The predictor
248 variables considered in the model selection process comprised *body mass*, *ground foraging*, *ground*
249 *nesting*, *preferred habitat*, aggregation at *waterholes*, use of *urban* areas and *island-endemicity*
250 (Table 2: italics indicate name used in reporting of modelling results). We log-transformed body

251 mass and allowed the effect of body mass to be non-linear by introducing a quadratic term,
252 stipulating its inclusion in a model only with the linear term. All continuous variables were
253 standardised by dividing by two times the standard deviation (Gelman 2008).

254

255 To consider model uncertainty, we took a model averaging approach to the analysis which
256 incorporates estimates from multiple candidate models weighted according to Akaike Information
257 Criterion with correction for small sample size (AIC_c) (Burnham and Anderson 2002). In this way, we
258 examined several competing models simultaneously to identify the top set of models (95%
259 confidence model set). These top models were averaged to obtain parameter estimates and
260 predictions were generated based on full model-averaged coefficients obtained from summed
261 Akaike weight (R package MuMIn: Barton 2016). The *abundance*-distribution factor (Table 2) was
262 used as an offset variable, and stipulated *a priori* for inclusion in all candidate models.

263

264 To identify a single optimal model for the purpose of visualisation of variable effects (R package
265 visreg: Breheny and Burchett 2016), relative variable importance values (w_+), defined as the sum of
266 Akaike weights for all models containing a given predictor variable, were used to identify only highly
267 influential variables ($w_+ \geq 0.73$, equivalent to an AIC difference of 2 which is widely used to assess a
268 'clear' effect: Richards (2005)) for inclusion in the optimal model. Optimal model validation was
269 conducted by calculation of Variance Inflation Factors (VIF) (car package: Fox and Weisberg 2011) to
270 test for multicollinearity among predictor variables, the dispersion statistic to test the fit of the
271 distribution, Cook's distances to check for observations with disproportionately high influence, and
272 adjusted McFadden Pseudo R² (pscl package: Jackman 2015) to estimate the deviance explained by
273 the model. Pearson residuals were plotted against fitted values, as well as included and excluded
274 covariates, to check for homogeneity, independence and model fit. For categorical variables with
275 more than two levels (i.e. *preferred habitat* and *ground nesting*), we used the 'glht' function (R
276 package multcomp: Hothorn et al. 2008) to identify significant differences among categories.

277

278 To answer the question: 'what is the relative likelihood that a cat will prey upon a bird species?'
279 predictions (P_{cat}) were generated by offsetting the abundance variable: for this question, bird species
280 that are more common are likely to rank highly. To answer the question 'among all bird species,
281 what is the relative likelihood of an individual bird being killed by a cat?' abundance was held
282 constant at the mean when generating predictions (P_{bird}). This question relates to a bird species'
283 relative *per capita* rate of predation by cats – for example, a rare species for which 20% of
284 individuals are killed by cats per year would rank higher than a very common species for which only
285 10% of individuals are killed by cats per year. This prediction is the likelihood of an individual of a
286 bird species being killed by cats (relative to all other bird species), given its ecological and other
287 traits. Note that it is not an explicit probability of an individual of that bird species being killed by
288 cats over any particular time period.

289

290 The modelling was repeated with the dependent variable being the number of separate sources
291 reporting cat-predation (rather than whether or not there were any cat-predation records for a bird
292 species in our compilation). The same predictor variables as used above in the binary analysis (Table
293 2) were considered in the model selection process for number of sources. To model this count data
294 we used negative binomial GLM's and predictions were generated from model-averaged coefficients
295 obtained from a top 95% confidence model set (R package MuMIn: Barton 2016). This parallel

296 analysis recognises that there are somewhat different biases in each approach: for example, use of
297 only presence/absence of predation records treats a bird species that may have had only a single
298 and unusual record of cat predation as equivalent to a species with numerous records indicating cat
299 predation on that species occurs frequently; whereas use of number of sources reinforces the bias
300 that species that are common, much-studied and occur in areas overlapping human population
301 centres are likely to be more frequently recorded as cat-predated, even if the incidence of such cat
302 predation is actually comparable to rare species occurring in remote areas.

303

304

305 **Results**

306

307 *Collation*

308 We collated records in Australia of 339 native bird species (of which one species was a vagrant to
309 Australia), with this tally comprising 45.6% of the 741 Australian native bird species, excluding
310 vagrants (Appendix B). Cat predation was also stated in our sources as presumed or implied in
311 Australia, or reported elsewhere, for a further 56 native bird species (of which three are vagrants to
312 Australia). Our compilation also includes 18 introduced bird species reported as killed by cats in
313 Australia (Appendix B). These tallies represent major advances from the previous compilations, of
314 cat predation records on 113 Australian native bird species reported by Doherty et al. (2015), and on
315 186 bird species (native and introduced) reported by (Paton 1990; Paton 1991; Paton 1993)

316

317 Our compilation includes records of cat predation in Australia for 75 bird species listed as extinct or
318 threatened by the IUCN or (with one or more subspecies listed as threatened) under Australia's
319 *Environment Protection and Biodiversity Conservation (EPBC) Act 1999* (Appendix B). This includes
320 one extinct species (Paradise Parrot *Psephotus pulcherrimus*), and 23 species listed as threatened by
321 the IUCN (40% of the 58 IUCN-listed threatened bird species occurring (other than as vagrants) in
322 Australia). Our collation includes records of cat predation for 71 of the 117 bird species (i.e. 61%)
323 that are listed as threatened species under the EPBC Act or have one or more subspecies so listed.
324 Again, these tallies represent major advances from the previous compilations, notably of cat
325 predation on two threatened Australian bird species recorded by Doherty et al. (2015).

326

327 Bird species reported to be killed by cats in Doherty et al. (2015) were more widespread and
328 abundant (mean 6624 Atlas records per species; s.e. 768), and more likely to have been well-studied
329 (mean of 11084 individuals banded; s.e. 2466), than the additional bird species recorded as killed by
330 cats in the current compilation (mean of 2247 Atlas records (s.e. 253); mean of 4234 individuals
331 banded (s.e. 1182)): i.e. our inclusion of more diverse sources served to capture cat-predation
332 records of more rare and restricted bird species than the previous compilation. However, both sets
333 of species were also more widespread and common, and more likely to have been studied, than bird
334 species for which we could locate no records of being killed by cats (mean of 895 Atlas records (s.e.
335 100); mean of 1988 individuals banded (s.e. 346)). The differences among these three groups of
336 species (i.e. recorded in previous compilation as cat-predated, newly recorded here as cat-predated,
337 or with no records of cat predation) were highly significant ($H=119.1$, $p<0.0001$ for Atlas records;
338 $H=93.7$, $p<0.0001$ for numbers of birds banded).

339

340 *Modelling*

341 A 95% confidence set of logistic regression models for extant native birds generated eight models
342 from summed AIC_c weights. All predictor variables other than *urban* and *waterholes* were highly
343 influential (Tables 3, 4).

344

345 For the optimal model containing only highly influential variables, VIF was less than 1.3, suggesting
346 that any collinearity among variables was unlikely to affect statistical inference (Zuur et al. 2010).
347 Further model validation techniques confirmed no dispersion issues, Cook's distances were <0.1,
348 residuals were unbiased and homoscedastic, and adjusted McFadden Pseudo R² of 0.16 indicated
349 good model fit (McFadden 1974).

350

351 From the optimal model (Akaike weight $w_i = 0.35$), the relative likelihood of a bird species being
352 preyed upon by cats was higher for bird species that forage on the ground, are of medium size (ca.
353 60-300 g) and are island endemics. Bird species that nest in Australia on the ground were more likely
354 to be preyed upon by cats than were bird species that were non-breeding visitors ($p < 0.001$), and
355 those that typically nest in Australia >1 m above ground ($p < 0.01$) (Fig. 1). *Preferred habitat* was also
356 associated with likelihood of being killed by a cat, with bird species primarily occurring in
357 rainforests/mangroves being less likely to be killed by cats than those associated with grasslands
358 ($p < 0.05$), shrublands/heathlands ($p < 0.05$), and open forests/woodlands ($p < 0.001$); coastal/marine
359 bird species also had a relatively high likelihood of being killed by cats.

360

361 Based on modelling of traits, the bird species that cats are most likely to prey upon are mostly
362 widespread and common species that forage (and/or nest) on or near the ground. These include
363 species such as Masked Lapwing *Vanellus miles*, Australasian Pipit *Anthus novaeseelandiae*, Superb
364 Fairy-wren *Malurus cyaneus*, Common Blackbird *Turdus merula*, Silver Gull *Chroicocephalus*
365 *novaehollandiae*, Yellow-rumped Thornbill *Acanthiza chrysorrhoa* and Striated Pardalote *Pardalotus*
366 *striatus* (Appendix D).

367

368 When bird abundance is held constant to provide predictions of the *per capita* likelihood of a bird
369 species being preyed upon by cats, the ordering of bird species is very different (Appendix E).
370 Reflecting the strong influence of the *island*-endemicity variable in the models, the species with
371 highest predicted *per capita* likelihood of being preyed upon by cats are island endemic species,
372 including several that are now extinct. With the small set of island-endemic species and the island
373 variable excluded, the 40 bird species with highest predicted *per capita* likelihood of being preyed
374 upon by cats is listed in Table 5, and values for all species given in Appendix F. Species with highest
375 modelled *per capita* likelihood of cat predation included many relatively localised, uncommon and
376 little-studied species, with some consistent groupings, notably of quail-thrush *Cinlosoma* spp.,
377 button-quail *Turnix* spp., and some ground-dwelling pigeons.

378

379 Results from modelling that used, as the dependent variable, the number of documented sources of
380 cat predation per bird species were consistent with modelling using only presence/absence of cat
381 predation records: detailed results are presented in Appendix G.

382

383

384 **Discussion**

385

386 Our compilation greatly increases the number of Australian bird species, and number of threatened
387 bird species, known to be preyed upon by cats. This is largely because we use a far more diverse set
388 of primary sources than the previous national compilation (Doherty et al. 2015), whose sources were
389 largely restricted to studies that focused on feral cat diet (rather than also including reports of
390 factors involved in bird mortality) and hence tended to include mostly common and widespread bird
391 species. Notwithstanding our extensive search of the literature, our results also indicate that our
392 compilation may retain some bias against recording predation by cats on less common and more
393 localised bird species.

394

395 Our results are largely consistent with previous studies that have reported that a very broad range of
396 bird species are preyed upon by cats, and that particular traits render some bird species more
397 susceptible to such predation (Dickman 1996; Kutt 2012; Lepczyk et al. 2004; Paltridge et al. 1997).
398 Our models indicate that predation by cats is most likely for bird species that nest and forage on the
399 ground and occur mostly in relatively open habitats (rather than rainforests and mangroves). A
400 preference by cats for bird species that forage and/or nest on the ground has been reported
401 previously in Australia (Paltridge et al. 1997; Paton 1991) and elsewhere (Dunn and Tessaglia 1994;
402 Lepczyk et al. 2004; Mead 1982).

403

404 Our demonstration that bird species' preferred habitat also influences the likelihood of a bird
405 species being preyed upon by cats may have several explanations. Our analysis may not have
406 completely overcome marked unevenness in the information base arising because there have been
407 few studies of the ecology and diet of cats in rainforest and mangrove habitats (Doherty et al. 2015).
408 The relative lack of such studies may itself be because these comprise only a small proportion of
409 Australia's continental area. To some extent, this bias can be redressed through information derived
410 from autecological studies of bird species associated with these closed forest habitats. Although
411 there are notable autecological studies of some Australian rainforest and mangrove bird species
412 (Frith and Frith 1995; Heinsohn et al. 2009; Laurance and Grant 1994; Noske 1996, 2001), few report
413 rates and causes of mortality. The relatively low likelihood of predation by cats predicted here for
414 bird species associated with rainforests and mangroves may be real rather than an artefact of
415 sampling unevenness. The likelihood of cat predation on birds is probably low in closed forest
416 environments because cat density is relatively low in such environments (Legge et al. 2017), and/or
417 because characteristics of the understorey of these environments may reduce cat hunting efficiency,
418 and/or because many bird species in these environments are canopy-dwellers. We cannot readily
419 partition the relative influence of these potential explanations, and more research on the abundance
420 and impacts of cats in these environments is warranted. The relatively higher likelihood of cat
421 depredation for birds occurring in coastal/marine habitats than for birds in freshwater wetland
422 habitats is probably because cats kill many seabirds that nest colonially on land, whereas most
423 freshwater wetland birds are offered some protection from cat predation by the water itself.

424

425 We also demonstrate that predation by cats is most likely for bird species of intermediate body mass
426 (ca. 60-300 g). Bird body size has been linked with likelihood of cat predation in previous studies: for
427 example, in north-eastern Queensland, Kutt (2012) found that cat predation was selective for birds
428 in the 10-50 g range, Dickman (1996) proposed that feral cats on the Australian mainland prefer
429 birds up to 200 g, and Paton (1991) considered that most birds taken in urban and peri-urban areas
430 of south-eastern Australia were <100 g. Cats' preferred bird size range may be difficult to

431 circumscribe neatly, given that the presence of large birds in cat diets may represent consumption of
432 their carrion or take of chicks or eggs.

433

434 We found no indication that the likelihood of cat predation was higher for bird species that
435 aggregate at water sources, in contrast to such preference being reported for some studies in arid
436 Australia (Paltridge et al. 1997). This difference may be because our assessment was continental in
437 scope, and aggregations of bird species at water sources are largely a phenomenon of arid and semi-
438 arid areas.

439

440 Although there were relatively few island-endemic bird species in our data set (24 species), these
441 few island-endemic species are strongly associated with relatively high predation risk, with the 18
442 bird species with highest predicted likelihood of predation by cats all being island-endemic species
443 (Appendix E). The susceptibility of island-endemic species, including bird species, to be killed by cats
444 is well established, with island endemic bird species contributing disproportionately to all known
445 bird extinctions, in large part due to introduced cats (Blackburn et al. 2005; Doherty et al. 2016;
446 Medina et al. 2011; Nogales et al. 2004). Furthermore, where cats are present on Australian islands,
447 their densities are, on average, an order of magnitude higher than on comparable areas of the
448 mainland (Legge et al. 2017), and such elevated densities of cats could contribute to the greater
449 likelihood of island birds being killed by cats. Furthermore, cats on Australian islands typically
450 consume a higher proportion of birds in their diet than do cats in comparable mainland areas
451 (Doherty et al. 2015; Woinarski et al. in press).

452

453 The models allowed us to estimate the likelihood of predation by cats for every Australian bird
454 species (Table 5; Appendices E, F), with control of many biases in our documentation. To our
455 knowledge, there are no comparable estimates of predation risk for entire continental bird faunas
456 elsewhere. These predicted values provide a general indication of the types of birds that may be
457 most detrimentally affected by cat predation, with high *per capita* likelihood of cat predation
458 particularly for island endemic, ground-nesting, ground-foraging and medium-sized species. Given
459 their high predicted rates of *per capita* cat-predation, we consider there may be particular cause for
460 conservation attention for all island-endemic bird species, ground-dwelling pigeons and doves
461 (*Phaps*, *Petrophassa*, *Geophaps* spp.), quail-thrush *Cinclosoma* spp., quail *Coturnix* spp., Plains-
462 Wanderer *Pedionomus torquatus* and button-quail *Turnix* spp (Table 5). Appropriate management
463 responses may include as enhanced management of cats in areas important for these bird species,
464 monitoring of population trends for these species and autecological studies. Although some of these
465 species are recognised to be of conservation concern, many have not hitherto been considered as
466 meriting particular conservation attention.

467

468 Our models included only a small number of traits, and some of these were greatly simplified from
469 original sources, so we may well have lost much of their ecological nuance. Our models also did not
470 include some traits (such as conspicuousness of plumage, and wariness) that may influence the
471 likelihood of a bird species being preyed upon by cats but were not readily parameterised. Although
472 challenging to parameterise, inclusion of appropriate measures for these characteristics could in
473 future help refine our modelling and improve its predictive power.

474

475 Records of predation by cats, or the predicted likelihoods of such predation, do not necessarily
476 correspond to conservation impact or consequences to the population viability of any bird species.
477 Impacts may also be influenced by the relative abundance of a bird species, the relative abundance
478 of cats, the relative availability of other prey to cats, a bird species' reproductive output and life
479 history, the array of other threats, and the interaction of other factors (such as fire regime, habitat
480 fragmentation and livestock grazing) that may serve to increase or decrease the severity of
481 predation impacts (Graham et al. 2013; Leahy et al. 2015; McGregor et al. 2015; McGregor et al.
482 2014; McGregor et al. 2016). Notwithstanding these caveats, the predicted values reported here of
483 relative *per capita* likelihood of being killed by cats for every Australian bird species are probably
484 more robust indicators of the potential threat of cat depredation to individual bird species than is a
485 simple documentation of whether or not there are predation records reported.

486

487 Given the now near-pervasiveness of cats across the Australian landscape, including many islands
488 and almost all conservation reserves (Legge et al. 2017), and that cats kill on average ca. 377 million
489 Australian birds per year (Woinarski et al. in press), our demonstration here that many more
490 Australian bird species (particularly threatened species) are preyed upon by cats than previously
491 recognised suggests that there is an urgent need to undertake more intensive studies of the impacts
492 of cat predation on the population viability of at least those bird species most likely to be
493 susceptible. Our results also support recent management initiatives to increase the currently very
494 small proportion of Australia that is free of cats (either on islands or within fenced predator-
495 enclosures) and the area in which cats are intensively controlled (Commonwealth of Australia 2015).

496

497

498 **Acknowledgements**

499

500 The data collation, analysis and preparation of this paper was supported by the Australian
501 Government's National Environmental Science Programme (Threatened Species Recovery Hub). We
502 thank David Drynan and the Australian Bird & Bat Banding Scheme (Department of the Environment
503 and Energy) for provision of tallies of cat-killed birds in bird-banding records. We thank the Museum
504 and Art Gallery of the Northern Territory (and curator Gavin Dally), Museum of Victoria (Laura Cook),
505 Tasmanian Museum and Art Gallery (Belinda Bauer), Western Australian Museum (Rebecca Bray),
506 Australian National Wildlife Collection (CSIRO: Leo Joseph), Queensland Museum (Heather Janetzki,
507 Andrew Amey), South Australian Museum (David Stemmer, Philippa Horton), and Australian
508 Museum (Cameron Slatyer, Mark Eldridge) for records of birds in their collection reported as cat-
509 killed. We thank the Australian Research Council for grant funding (project DP 140104621) to CRD.
510 This paper rests on data arising from the dedicated labours of many people who have searched for
511 and through cat faeces and the digestive tracts of dead cats: that effort is much appreciated. We
512 also thank two anonymous referees for comments that improved this paper.

513

514

515 **References**

516

517 Abbott, I., 2008. The spread of the cat, *Felis catus*, in Australia: re-examination of the current
518 conceptual model with additional information. Conservation Science Western Australia 7, 1-17.

519 Barton, K., 2016. MuMIn: Multi-Model Inference. R package version 1.15.6. [https://CRAN.R-](https://CRAN.R-project.org/package=MuMIn)
520 [project.org/package=MuMIn](https://CRAN.R-project.org/package=MuMIn).

521 Blackburn, T.M., Cassey, P., Duncan, R.P., Evans, K.L., Gaston, K.J., 2004. Avian extinction and
522 mammalian introductions on oceanic islands. *Science* 305, 1955-1958.

523 Blackburn, T.M., Petchey, O.L., Cassey, P., Gaston, K.J., 2005. Functional diversity of mammalian
524 predators and extinction in island birds. *Ecology* 86, 2916-2923.

525 Blancher, P., 2013. Estimated number of birds killed by house cats (*Felis catus*) in Canada. *Avian*
526 *Conservation and Ecology* 8, 3.

527 Bonnaud, E., Medina, F.M., Vidal, E., Nogales, M., Tershy, B., Zavaleta, E., Donlan, C.J., Keitt, B., Le
528 Corre, M., Horwath, S.V., 2011. The diet of feral cats on islands: a review and a call for more studies.
529 *Biological Invasions* 13, 581-603.

530 Bradshaw, J.W.S., 1992. The behaviour of the domestic cat. CAB International, Wallingford UK.

531 Breheny, P., Burchett, W., 2016. visreg: Visualization of Regression Models. R package version 2.3-0.
532 <https://CRAN.R-project.org/package=visreg>.

533 Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical
534 information-theoretic approach. Springer, Berlin.

535 Commonwealth of Australia, 2015. Threatened species strategy, Canberra.

536 Dauphiné, N., Cooper, R.J., 2009. Impacts of free-ranging domestic cats (*Felis catus*) on birds in the
537 United States: a review of recent research with conservation and management recommendations, In
538 Tundra to tropics: connecting birds, habitats and people. Proceedings of the 4th International
539 Partners in Flight Conference, 13-16 February 2008, McAllen, Texas. eds T.D. Rich, C. Arizmendi,
540 D.W. Demarest, C. Thompson, pp. 205-219. Partners in Flight, McAllen, Texas.

541 Department of the Environment, 2015. Threat abatement plan for predation by feral cats, Canberra.

542 Dickman, C.R., 1996. Overview of the Impacts of Feral Cats on Australian Native Fauna. Australian
543 Nature Conservation Agency, Canberra.

544 Doherty, T.S., Davis, R.A., Etten, E.J.B., Algar, D., Collier, N., Dickman, C.R., Edwards, G., Masters, P.,
545 Palmer, R., Robinson, S., 2015. A continental-scale analysis of feral cat diet in Australia. *Journal of*
546 *Biogeography* 42, 964-975.

547 Doherty, T.S., Glen, A.S., Nimmo, D.G., Ritchie, E.G., Dickman, C.R., 2016. Invasive predators and
548 global biodiversity loss. *Proceedings of the National Academy of Sciences* 113, 11261–11265.

549 Dowling, B., Seebeck, J.H., Lowe, K.W., 1994. Cats and wildlife: results of a survey of wildlife
550 admitted for care to shelters and animal welfare agencies in Victoria. Department of Conservation
551 and Natural Resources, Melbourne.

552 Dunn, E.H., Tessaglia, D.L., 1994. Predation of birds at feeders in winter. *Journal of Field Ornithology*
553 65, 8-16.

554 Fancourt, B.A., 2015. Making a killing: photographic evidence of predation of a Tasmanian
555 pademelon (*Thylogale billardierii*) by a feral cat (*Felis catus*). *Australian Mammalogy* 37, 120-124.

556 Fox, J., Weisberg, S., 2011. An R Companion to Applied Regression, 2nd edn. Sage, Thousand Oaks
557 California.

558 Frith, C.B., Frith, D.W., 1995. Court site constancy, dispersion, male survival and court ownership in
559 the male Tooth-billed Bowerbird, *Scenopoeetes dentirostris* (Ptilonorhynchidae). *Emu* 95, 84-98.

560 Garnett, S.T., Duursma, D.E., Ehmke, G., Guay, P.-J., Stewart, A., Szabo, J.K., Weston, M.A., Bennett,
561 S., Crowley, G.M., Drynan, D., Dutson, G., Fitzherbert, K., Franklin, D.C., 2015. Biological, ecological,
562 conservation and legal information for all species and subspecies of Australian bird. *Scientific data* 2.
563 Gelman, A., 2008. Scaling regression inputs by dividing by two standard deviations. *Statistical*
564 *Methods* 27, 2865-2873.

565 Graham, C.A., Maron, M., McAlpine, C.A., 2013. Influence of landscape structure on invasive
566 predators: feral cats and red foxes in the brigalow landscapes, Queensland, Australia. *Wildlife*
567 *Research* 39, 661-676.

568 Heinsohn, R., Zeriga, T., Murphy, S., Igag, P., Legge, S., Mack, A.L., 2009. Do Palm Cockatoos
569 (*Probosciger aterrimus*) have long enough lifespans to support their low reproductive success? Emu
570 109, 183-191.

571 Higgins, P.J. ed., 1999. Handbook of Australian, New Zealand & Antarctic Birds. Volume 4. Parrots to
572 Dollarbirds. Oxford University Press, Melbourne.

573 Higgins, P.J., Davies, S.J.J.F. eds., 1996. Handbook of Australian, New Zealand & Antarctic Birds.
574 Volume 3. Snipe to Pigeons. Oxford University Press, Melbourne.

575 Higgins, P.J., Peter, J.M. eds., 2002. Handbook of Australian, New Zealand & Antarctic Birds. Volume
576 6. Pardalotes to Shrike-thrushes. Oxford University Press, Melbourne.

577 Higgins, P.J., Peter, J.M., Cowling, S.J. eds., 2006. Handbook of Australian, New Zealand & Antarctic
578 Birds. Volume 7. Boatbills to Starlings. Oxford University Press, Melbourne.

579 Higgins, P.J., Peter, J.M., Steele, W.K. eds., 2001. Handbook of Australian, New Zealand & Antarctic
580 Birds. Volume 5. Tyrant-flycatchers to Chats. Oxford University Press, Melbourne.

581 Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric models.
582 Biometrical Journal 50, 346-363.

583 Jackman, S., 2015. pscl: Classes and Methods for R Developed in the Political Science Computational
584 Laboratory, Stanford University. R package version 1.4.9. Department of Political Science, Stanford
585 University, Stanford.

586 Kutt, A.S., 2012. Feral cat (*Felis catus*) prey size and selectivity in north-eastern Australia:
587 implications for mammal conservation. Journal of Zoology 287, 292-300.

588 Laurance, W.F., Grant, J.D., 1994. Photographic identification of ground-nest predators in Australian
589 tropical rainforest. Wildlife Research 21, 241-247.

590 Leahy, L., Legge, S.M., Tuft, K., McGregor, H., Barmuta, L., Jones, M.E., Johnson, C.N., 2015. Amplified
591 predation after fire suppresses rodent populations in Australia's tropical savannas. Wildlife Research
592 42, 705-716.

593 Legge, S., Murphy, B.P., McGregor, H., Woinarski, J.C.Z., Augusteyn, J., Ballard, G., Baseler, M.,
594 Buckmaster, T., Dickman, C.R., Doherty, T., Edwards, G., Eyre, T., Fancourt, B., Ferguson, D., Forsyth,
595 D.M., Geary, W.L., Gentle, M., Gillespie, G., Greenwood, L., Hohnen, R., Hume, S., Johnson, C.N.,
596 Maxwell, N., McDonald, P., Morris, K., Moseby, K., Newsome, T., Nimmo, D., Paltridge, R., Ramsey,
597 D., Read, J., Rendall, A., Rich, M., Ritchie, E., Rowland, J., Short, J., Stokeld, D., Sutherland, D.R.,
598 Wayne, A.F., Woodford, L., Zewe, F., 2017. Enumerating a continental-scale threat: how many feral
599 cats are in Australia? Biological Conservation 206, 293-303.

600 Lepczyk, C.A., Mertig, A.G., Liu, J., 2004. Landowners and cat predation across rural-to-urban
601 landscapes. Biological Conservation 115, 191-201.

602 Loss, S.R., Will, T., Marra, P.P., 2012. Direct human-caused mortality of birds: improving
603 quantification of magnitude and assessment of population impact. Frontiers in Ecology and the
604 Environment 10, 357-364.

605 Loss, S.R., Will, T., Marra, P.P., 2013. The impact of free-ranging domestic cats on wildlife of the
606 United States. Nature Communications 4, 1396.

607 Loss, S.R., Will, T., Marra, P.P., 2015. Direct mortality of birds from anthropogenic causes. Annual
608 review of ecology, evolution, and systematics 46, 99-120.

609 Marchant, S., Higgins, P.J. eds., 1990. Handbook of Australian, New Zealand & Antarctic Birds.
610 Volume 1. Ratites to Ducks. Oxford University Press, Melbourne.

611 Marchant, S., Higgins, P.J. eds., 1993. Handbook of Australian, New Zealand & Antarctic Birds.
612 Volume 2. Raptors to Lapwings. Oxford University Press, Melbourne.

613 Mattingley, A.H.E., 1918. The ground-parrot (*Pezoporus formosus*). Emu 17, 216-218.

614 McFadden, D., 1974. Conditional logit analysis of qualitative choice behavior, In Frontiers in
615 Econometrics (Economic theory and mathematical economics). ed. P. Zarembka, pp. 105-142.
616 Academic Press, New York.

617 McGregor, H., Legge, S., Jones, M.E., Johnson, C.N., 2015. Feral cats are better killers in open
618 habitats, revealed by animal-borne video. PLoS ONE 10, e0133915.

619 McGregor, H.W., Legge, S., Jones, M.E., Johnson, C.N., 2014. Landscape management of fire and
620 grazing regimes alters the fine-scale habitat utilisation by feral cats. PLoS ONE 9, e109097.

621 McGregor, H.W., Legge, S.M., Jones, M.E., Johnson, C.N., 2016. Extraterritorial hunting expeditions
622 to intense fire scars by feral cats. Scientific Reports 6, 22559.

623 Mead, C.J., 1982. Ringed birds killed by cats. Mammal Review 12, 183-186.

624 Medina, F.M., Bonnaud, E., Vidal, E., Tershy, B.R., Zavaleta, E.S., Josh Donlan, C., Keitt, B.S., Corre,
625 M., Horwath, S.V., Nogales, M., 2011. A global review of the impacts of invasive cats on island
626 endangered vertebrates. Global Change Biology 17, 3503-3510.

627 Molsher, R.L., Newsome, A.E., Newsome, T.M., Dickman, C.R., 2017. Mesopredator management:
628 effects of red fox control on the abundance, diet and use of space by feral cats. PLoS ONE 12,
629 e168460.

630 Nogales, M., Martín, A., Tershy, B.R., Donlan, C., Veitch, D., Puerta, N., Wood, B., Alonso, J., 2004. A
631 review of feral cat eradication on islands. Conservation Biology 18, 310-319.

632 Nogales, M., Vidal, E., Medina, F.M., Bonnaud, E., Tershy, B.R., Campbell, K.J., Zavaleta, E.S., 2013.
633 Feral cats and biodiversity conservation: the urgent prioritization of island management. Bioscience
634 63, 804-810.

635 Noske, R.A., 1996. Abundance, zonation and foraging ecology of birds in mangroves of Darwin
636 Harbour, Northern Territory. Wildlife Research 23, 443-474.

637 Noske, R.A., 2001. The breeding biology of the mangrove gerygone, *Gerygone laevigaster*, in the
638 Darwin region, with notes on brood parasitism by the little bronze-cuckoo, *Chrysococcyx minutillus*.
639 Emu 101, 129-135.

640 Paltridge, R., Gibson, D., Edwards, G., 1997. Diet of the feral cat (*Felis catus*) in central Australia.
641 Wildlife Research 24, 67-76.

642 Paton, D., 1990. Domestic cats and wildlife: results from initial questionnaire. Bird Observer 696, 34-
643 35.

644 Paton, D.C., 1991. Loss of wildlife to domestic cats, In The impact of cats on native wildlife. ed. C.
645 Potter, pp. 64-69. Australian National Parks & Wildlife Service, Canberra.

646 Paton, D.C., 1993. Impacts of domestic and feral cats on wildlife, In Cat Management Workshop
647 Proceedings – 1993. eds G. Siepen, C. Owens, pp. 9-15. Queensland Department of Environment and
648 Heritage, Brisbane.

649 R Core Team, 2016. R: A language and environment for statistical computing. R Foundation for
650 Statistical Computing, Vienna.

651 Richards, S.A., 2005. Testing ecological theory using the information-theoretic approach: examples
652 and cautionary results. Ecology 86, 2805–2814.

653 Smith, G.T., Saunders, D.A., 1986. Clutch size and productivity in three sympatric species of cockatoo
654 (Psittaciformes) in the southwest of Western Australia. Australian Wildlife Research 13, 275-285.

655 Turner, D.C., Meister, O., 1988. Hunting behaviour of the domestic cat, In The domestic cat: the
656 biology of its behaviour. eds D.C. Turner, P. Bateson, pp. 111-121. Cambridge University Press,
657 Cambridge.

658 Woinarski, J.C.Z., Burbidge, A.A., Harrison, P.L., 2015. The ongoing unravelling of a continental fauna:
659 decline and extinction of Australian mammals since European settlement. Proceedings of the
660 National Academy of Sciences 15, 4531-4540.

661 Woinarski, J.C.Z., Murphy, B.P., Legge, S.M., Garnett, S.T., Lawes, M.J., Comer, S., Dickman, C.R.,
662 Doherty, T.S., Edwards, G., Nankivell, A., Paton, D., Palmer, R., Woolley, L.A., in press. How many
663 birds are killed by cats in Australia? Biological Conservation.

664 Zuur, A.F., Ieno, E.N., Elphick, C.S., 2010. A protocol for data exploration to avoid common statistical
665 problems. Methods in Ecology and Evolution 1, 3-14.

666

667

668

669
670
671
672

Table 1. Real or potential biases in documentation of records of cat predation, and constraints on modelling.

Potential bias	Response in this study to reduce bias
Studies of cat-predation will tend to report records of predation of more common and widespread bird species, and those occurring in areas in and around human population centres	We included information on predation from many diverse sources, including autecological studies of birds, rather than simply collations of cat diet; our modelling includes an offset for abundance, to allow derivation of a <i>per capita</i> estimate of predation risk
Observations of cat predation on birds will be biased towards larger and more distinctive birds	We included information on predation from many diverse sources, including autecological studies of birds, rather than simply collations of cat diet. The bias due to some bird species being more conspicuous or more easily identified mostly relates to the minority of records here that derive from pet-owners' reports
Observations of cat predation on birds will be biased towards bird species that have been the subject of intensive autecological studies	This bias was not entirely circumvented in our compilation or modelling. However, there are relatively few autecological studies of Australian bird species that include documentation of different sources of mortality, and our compilation used very diverse sources in addition to reports from autecological studies.
There have been relatively few studies of birds or cats in mangroves and rainforest habitats.	This bias was not entirely circumvented in our compilation or modelling, but other studies (Legge et al. 2017) indicate that cat density is likely to be relatively low in closed forest habitats.
There will be fewer records of cat predation on birds that became extinct soon after European settlement	This bias was not entirely circumvented in our compilation or modelling, but modelling indicated high predation risk for many extinct bird species anyway
Eggs and nestlings will be under-represented in samples because these may be quickly digested and unidentifiable in cat samples	This bias was not entirely circumvented in our compilation or modelling, but is unlikely to introduce any systematic bias for or against particular bird species
Larger birds may be included in cat samples but these may represent carrion rather than predation	This bias was not circumvented in our compilation or modelling, but our inclusion of predation information arising from assessments of causes of mortality within autecological studies of birds may redress this concern
Cats may kill birds but not consume them ('surplus kill'), and these killed birds will not be present in dietary samples	This factor should not introduce any major bias among bird species – i.e. although colonial bird species may be more likely to be

Consumption of a single individual of a large bird species may satiate cats, whereas it may require many small birds to satiate cats

subject to 'surplus killing' this should not affect our analysis, which is based on any records of bird species being killed rather than the tally of numbers of individuals being killed

Not a bias *per se* – simply recognises that more individuals of smaller bird species may be taken by cats than of larger birds

674 Table 2. Bird traits used in modelling. Note that we also used information presented in Garnett et al.
 675 (2015) to categorise bird species as vagrant or not, extinct or extant, native or introduced, and
 676 threatened or not.
 677

Parameter	Coding	Source	Comment
<i>Body mass</i>	Adult body mass (g)	Garnett et al. (2015)	Note that cat-predation records may relate to predation on much smaller chicks, or eggs
<i>Preferred habitat</i>	Categorical (as either 1=grassland, 2=shrubland/heathland, 3=woodland/open forest, 4=rainforest/mangrove, 5=freshwater, or 6=coastal/marine)	Simplified from Garnett et al. (2015) (see Appendix C)	
<i>Urban use</i>	Categorical (as 0=not reported to use urban habitats; 1=reported to use urban habitats)	Garnett et al. (2015)	
<i>Island endemic</i>	Categorical (as 0=not endemic to islands, or 1=endemic to islands)	Garnett et al. (2015)	
<i>Waterholes</i>	Categorical (as 0=typically does not aggregate to drink at waterholes; 1=often aggregates to drink at waterholes)	Derived anew from information presented in HANZAB series	
<i>Abundance and distributional extent</i>	Continuous	Garnett et al. (2015)	This parameter was a log-transformed measure of the total number of observational records of a species in two Atlases of Australian Birds (1977 to 1981, and 1998 to 2001). Note that the Atlas index did not include any records from oceanic islands, and may have some bias towards species occurring mostly in or near areas of higher human population density.
<i>Extent of research effort</i>	Continuous	Australian Bird and Bat Banding Scheme	The number of individual birds banded (per species) was included in preliminary models as an indicator of study effort, but this variable included extremely high values for an idiosyncratic set of species, so was excluded from models described here
<i>Ground-foraging</i>	Continuous, varying from 0 (does not feed on the ground) to 3 (feeds entirely on the ground)	Simplified from Garnett et al. (2015) (see Appendix C)	
<i>Ground-nesting</i>	Categorical (as either 0=not nesting in Australia, 1= typically nesting in shrubs, trees or other sites >1 m above ground; or 2=typically nesting on the ground or within 1 m of it)	Simplified from Garnett et al. (2015) (see Appendix C)	

678
679
680
681
682
683
684
685
686

Table 3. Best candidate models (95% confidence model set) used to test the effects of predictor variables on records of cat-predation. AICc is the Akaike Information Criterion with correction for small sample size; ΔAICc is a measure of change in AICc relative to the best model; Akaike weight w_i is the probability that model i is the best model. When present in candidate models, body mass includes both linear and quadratic terms. All models include the offset for abundance. For definitions of variables see Table 2.

Model	ΔAICc	w_i
Ground foraging + ground nesting + habitat + island + body mass	0.00	0.35
Ground nesting + habitat + island + body mass	1.46	0.17
Ground foraging + ground nesting + habitat + island + body mass + waterholes	1.92	0.13
Ground foraging + ground nesting + habitat + island + body mass + urban	1.97	0.13
Ground nesting + habitat + island + body mass + waterholes	3.07	0.08
Ground nesting + habitat + island + body mass + urban	3.37	0.06
Ground foraging + ground nesting + habitat + island + body mass + waterholes + urban	3.91	0.05
Ground nesting + habitat + island + body mass + waterholes + urban	5.02	0.03

687
688

689
690
691
692
693

Table 4. Relative importance values (w_+) of predictor variables. For definitions of variables see Table 2.

Variable	w_+
Island	1.00
Habitat	1.00
Ground nesting	1.00
Body mass	1.00
Body mass ²	1.00
Ground foraging	0.76
Urban	0.29
Waterholes	0.27

694
695
696

697
698 Table 5. The 40 bird species with highest relative *per capita* likelihood of being killed by cats (P_{bird}).
699 These results derive from modelling, across all non-vagrant bird species, of the relationship between
700 presence/absence of cat-predation records and bird traits, with bird abundance kept constant, and
701 the small set of island-endemic species omitted. Values given in table are estimated value and 95%
702 confidence interval (95% CI). *species extinct in Australia; **threatened species, or at least one
703 subspecies listed as threatened

704

Common name	Scientific name	fit	95% CI
Spotted Quail-thrush**	<i>Cinclosoma punctatum</i>	0.794	(0.690-0.870)
Chestnut-backed Button-quail	<i>Turnix castanotus</i>	0.793	(0.689-0.868)
Painted Button-quail**	<i>Turnix varius</i>	0.792	(0.689-0.868)
Buff-breasted Button-quail**	<i>Turnix olivii</i>	0.792	(0.689-0.868)
White-quilled Rock-Pigeon	<i>Petrophassa albipennis</i>	0.792	(0.679-0.872)
Chestnut-quilled Rock-Pigeon	<i>Petrophassa rufipennis</i>	0.792	(0.678-0.873)
Partridge Pigeon**	<i>Geophaps smithii</i>	0.791	(0.677-0.872)
Brush Bronzewing	<i>Phaps elegans</i>	0.791	(0.676-0.872)
Squatter Pigeon**	<i>Geophaps scripta</i>	0.790	(0.675-0.872)
Chestnut Quail-thrush	<i>Cinclosoma castanotus</i>	0.789	(0.647-0.885)
Chestnut-breasted Quail-thrush	<i>Cinclosoma castaneothorax</i>	0.788	(0.646-0.883)
Sandstone Shrike-thrush	<i>Colluricincla woodwardi</i>	0.787	(0.685-0.863)
Red-backed Kingfisher	<i>Todiramphus pyrrhopygius</i>	0.787	(0.684-0.863)
Cinnamon Quail-thrush**	<i>Cinclosoma cinnamomeum</i>	0.786	(0.645-0.882)
Rufous Scrub-bird**	<i>Atrichornis rufescens</i>	0.784	(0.681-0.860)
Paradise Parrot*	<i>Psephotus pulcherrimus</i>	0.782	(0.672-0.863)
Southern Scrub-robin	<i>Drymodes brunneopygia</i>	0.776	(0.636-0.874)
Rufous Songlark	<i>Cinloramphus mathewsi</i>	0.776	(0.670-0.854)
Common Blackbird	<i>Turdus merula</i>	0.774	(0.673-0.851)
Western Ground Parrot**	<i>Pezoporus flaviventris</i>	0.772	(0.622-0.874)
Bush Stone-curlew	<i>Burhinus grallarius</i>	0.772	(0.651-0.860)
Eastern Ground Parrot	<i>Pezoporus wallicus</i>	0.771	(0.622-0.873)
Rufous Bristlebird**	<i>Dasyornis broadbenti</i>	0.769	(0.616-0.874)
Crested Bellbird	<i>Oreoica gutturalis</i>	0.768	(0.621-0.870)
California Quail	<i>Callipepla californica</i>	0.767	(0.650-0.853)
Stubble Quail	<i>Coturnix pectoralis</i>	0.767	(0.652-0.852)
Brown Quail	<i>Coturnix ypsilophora</i>	0.767	(0.652-0.852)
Banded Lapwing	<i>Vanellus tricolor</i>	0.766	(0.650-0.853)
Inland Dotterel	<i>Charadrius australis</i>	0.765	(0.651-0.850)
Night Parrot**	<i>Pezoporus occidentalis</i>	0.764	(0.641-0.855)
Spinifex Pigeon	<i>Geophaps plumifera</i>	0.764	(0.640-0.854)
Western Whipbird**	<i>Psophodes nigrogularis</i>	0.763	(0.618-0.865)
Eastern Bristlebird**	<i>Dasyornis brachypterus</i>	0.762	(0.618-0.865)
Plains-wanderer**	<i>Pedionomus torquatus</i>	0.762	(0.650-0.847)
Noisy Scrub-bird**	<i>Atrichornis clamosus</i>	0.762	(0.617-0.864)
Chirruping Wedgebill	<i>Psophodes cristatus</i>	0.761	(0.616-0.863)
Chiming Wedgebill	<i>Psophodes occidentalis</i>	0.760	(0.616-0.862)

Brown Songlark	<i>Cincloramphus cruralis</i>	0.760	(0.648-0.845)
Flock Bronzewing	<i>Phaps histrionica</i>	0.760	(0.634-0.852)
Rock Dove	<i>Columba livia</i>	0.760	(0.634-0.852)

705




Figure 1. Relationships between the *per capita* likelihood of being preyed upon by a cat (P_{bird}) and key predictor variables (while holding all other variables at fixed median levels (continuous variables) and most common category (categorical variables) and offsetting for bird species abundance by holding

abundance constant at the mean), derived from the optimal logistic regression model. Continuous lines represent fits to the model's predicted values and grey area indicates 95% confidence interval of model fits. Codes for categorical variables: island (0=not endemic to islands, 1=endemic to islands); ground nesting (0=does not breed in Australia, 1=nests >1 m above ground, 2=nests <1 m from ground); habitat (G=grassland, SH=shrubland/heathland, OF=woodland/open forest, RF=rainforest/mangrove, FW=freshwater, CM=coastal/marine).