
Charles Darwin University

Patterns of genetic structuring at the northern limits of the Australian smelt (*Retropinna semoni*) cryptic species complex

Islam, Md Rakeb Ul; Schmidt, Daniel J.; Crook, David A.; Hughes, Jane M.

Published in:
PEERJ

DOI:
[10.7717/peerj.4654](https://doi.org/10.7717/peerj.4654)

Published: 01/01/2018

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Islam, M. R. U., Schmidt, D. J., Crook, D. A., & Hughes, J. M. (2018). Patterns of genetic structuring at the northern limits of the Australian smelt (*Retropinna semoni*) cryptic species complex. *PEERJ*, 2018(5), [e4654]. <https://doi.org/10.7717/peerj.4654>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Patterns of genetic structuring at the northern limits of the Australian smelt (*Retropinna semoni*) cryptic species complex

Md Rakeb-Ul Islam^{1,*}, Daniel J. Schmidt^{1,*}, David A. Crook^{2,*} and Jane M. Hughes^{1,*}

¹ Australian Rivers Institute, Griffith University, Brisbane, Australia

² Research Institute for Environment and Livelihoods, Charles Darwin University, Darwin, NT, Australia

* These authors contributed equally to this work.

ABSTRACT

Freshwater fishes often exhibit high genetic population structure due to the prevalence of dispersal barriers (e.g., waterfalls) whereas population structure in diadromous fishes tends to be weaker and driven by natal homing behaviour and/or isolation by distance. The Australian smelt (Retropinnidae: *Retropinna semoni*) is a native fish with a broad distribution spanning inland and coastal drainages of south-eastern Australia. Previous studies have demonstrated variability in population genetic structure and movement behaviour (potamodromy, facultative diadromy, estuarine residence) across the southern part of its geographic range. Some of this variability may be explained by the existence of multiple cryptic species. Here, we examined genetic structure of populations towards the northern extent of the species' distribution, using ten microsatellite loci and sequences of the mitochondrial *cyt b* gene. We tested the hypothesis that genetic connectivity among rivers should be low due to a lack of dispersal via the marine environment, but high within rivers due to dispersal. We investigated populations corresponding with two putative cryptic species, SEQ-North (SEQ-N), and SEQ-South (SEQ-S) lineages occurring in south east Queensland drainages. These two groups formed monophyletic clades in the mtDNA gene tree and among river phylogeographic structure was also evident within each clade. In agreement with our hypothesis, highly significant overall F_{ST} values suggested that both groups exhibit very low dispersal among rivers (SEQ-S $F_{ST} = 0.13$; SEQ-N $F_{ST} = 0.27$). Microsatellite data indicated that connectivity among sites within rivers was also limited, suggesting dispersal may not homogenise populations at the within-river scale. Northern groups in the Australian smelt cryptic species complex exhibit comparatively higher among-river population structure and smaller geographic ranges than southern groups. These properties make northern Australian smelt populations potentially susceptible to future conservation threats, and we define eight genetically distinct management units along south east Queensland to guide future conservation management. The present findings at least can assist managers to plan for effective conservation and management of different fish species along coastal drainages of south east Queensland, Australia.

Submitted 25 September 2017

Accepted 31 March 2018

Published 3 May 2018

Corresponding author

Md Rakeb-Ul Islam,
r.islam@griffithuni.edu.au

Academic editor

Mohammad Shamsur Rahman

Additional Information and
Declarations can be found on
page 16

DOI 10.7717/peerj.4654

© Copyright
2018 Islam et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Aquaculture, Fisheries and Fish Science, Conservation Biology, Ecology, Genetics, Zoology

Keywords Facultative diadromy, Dispersal, Population structure, Cryptic species, Isolation by distance

INTRODUCTION

Genetic structure in aquatic fauna is strongly influenced by the characteristics of the ambient environment. Freshwater species typically exhibit higher levels of genetic differentiation than those living in estuarine or marine habitats (Ward, Woodwark & Skibinski, 1994; Sharma & Hughes, 2009). These greater levels of genetic structure in freshwater fish from different drainages are the result of the isolating nature of drainage systems and relatively smaller population sizes compared with marine species (McGlashan & Hughes, 2002; Huey, Baker & Hughes, 2010). Strong to moderate genetic structure is observed in many freshwater fish species suggesting restricted gene flow and limited dispersal among populations (Leclerc et al., 2008; Pereira, Foresti & Oliveira, 2009; Huey, Baker & Hughes, 2010). Movement by obligate freshwater organisms is limited to the water column and the freshwater environment, preventing inter-catchment dispersal via the sea (Burridge et al., 2008; Hughes, Schmidt & Finn, 2009; Bernays et al., 2015). In addition, various in-stream barriers to dispersal including waterfalls, dam walls, habitat heterogeneity, dried river reaches, and steep catchment gradients may act to restrict gene flow among populations within catchments (Amoros & Bornette, 2002; Huey, Baker & Hughes, 2010). As a consequence of the physical limitations to dispersal in freshwater environments, populations of aquatic organisms are often highly differentiated both among and within catchments (McGlashan & Hughes, 2000; Hughes, 2007; Sharma & Hughes, 2009). For a particular species, it is essential to understand the levels of population differentiation, genetic diversity, and rates of gene flow among populations for proper conservation and management of freshwater ecosystems (Geist, 2011).

The southern smelt (Retropinnidae: *Retropinna* spp.) is a common fish species distributed throughout the rivers of south-eastern Australia (McDowall, 1996). Individuals reach a maximum length of about 100 mm total length (TL), although adults are usually 50–60 mm TL (Pusey, Kennard & Arthington, 2004). Australian smelts are currently formally recognised as two described species *R. semoni* Weber, and *R. tasmanica* McCulloch, but recent genetic analyses have identified a complex of five or more cryptic species across their geographic range based on allozymes, microsatellites and mitochondrial DNA data (Hammer et al., 2007; Hughes et al., 2014; Schmidt, Islam & Hughes, 2016). Otolith chemistry studies in the southern part of their distribution have shown that Australian smelt exhibit a range of life history patterns, including freshwater residency, facultative diadromy and estuarine residency (Crook, Macdonald & Raadik, 2008; Hughes et al., 2014). In inland regions of Australia, large numbers of Australian smelt have been observed moving upstream through fishways (e.g., Baumgartner & Harris, 2007) and the species is widely described as potamodromous (i.e., migration within freshwater for the purpose of breeding) (e.g., Rolls, 2011). Nonetheless, Woods et al. (2010) found strong genetic structure among inland populations of Australian smelt and suggested low levels of dispersal in at

least some regions. These differences between studies could reflect differing life-histories among the cryptic species.

In most studies to date, diadromous behaviour has been shown to facilitate genetic connectivity among river catchments and typically results in “isolation-by-distance” (IBD) patterns of population genetic structure (Keenan, 1994; Jerry & Baverstock, 1998). In Australian smelt, however, there is strong genetic differentiation among catchments across the southern part of the range—even among populations containing diadromous individuals, suggesting high retention of fish within estuaries and a lack of marine dispersal (Hughes *et al.*, 2014). The aim of the current study was to examine patterns of genetic connectivity of populations in the north of the geographic range of Australian smelt, which contains two putative cryptic species (‘SEQ’, ‘CEQ’ *sensu.* Hammer *et al.*, 2007) that differ from those studied in detail previously (‘MTV’, ‘SEC’ *sensu.* This study focuses on the SEQ lineage which was further subdivided into northern and southern groups (SEQ-N, SEQ-S) following Page & Hughes (2014). Mitochondrial DNA sequence data combined with nuclear data from 10 microsatellite loci were used to test the hypotheses that, (i) northern *R. semoni* would display high population structure among rivers similar to southern populations; and (ii) that genetic structure within rivers would be low due to dispersal within rivers.

MATERIALS AND METHODS

Sampling strategy

A total of 389 individual samples were collected from 15 locations in south-east Queensland, Australia (Fig. 1; Table 1A). Samples were collected using a hand—held seine net from an upstream and a downstream site from each of eight river systems (except the Noosa River—downstream only). We aimed to collect at least 30 individuals per site but this was not always possible, as the species was not abundant in all rivers. Fin clips or entire individuals were placed in 95% ethanol in the field and stored prior to preparation for analysis. All procedures were carried out according to Australian Ethics Commission protocol number ENV/23/14/AEC.

Molecular methods

Genomic DNA was extracted from fin tissue using the DNeasy Blood and Tissue kits (Qiagen, Germantown, MD, USA) following the manufacturer’s directions. Microsatellite markers developed for *R. semoni* were amplified and genotyped using primers developed by Islam, Schmidt & Hughes (2017). Ten loci were screened across all individuals. The ten loci were BS18, BS3, BS4, BS5, BS20, BS21, BS22, BS24, BS8 and MS24. Microsatellite screening was carried out in 10 µl polymerase chain reaction (PCR) consisting of 0.5 µl of genomic DNA, 0.2 mM reverse primer, 0.05 mM tailed forward primer, 0.2 mM tailed fluorescent tag (either FAM, VIC, NED or PET, Applied Biosystems), 1 × PCR buffer (Astral Scientific, Caringbah, Australia) and 0.02 units of *taq* polymerase (Astral Scientific, Caringbah, Australia). The following basic thermocycler settings for PCR were performed: initial denaturation at 94 °C for 4 min, followed by 35 cycles at 94 °C for 1 min, 57 °C for 30 s, 72 °C for 1 min and a final extension at 72 °C for 7 min. Fluorescently




Figure 1 Localities where specimens of *R. semoni* were collected during this study in south-east Queensland, Australia. The green circle denotes the sampling site. See Tables 1A and 1B for site codes and locations.

Full-size  DOI: [10.7717/peerj.4654/fig-1](https://doi.org/10.7717/peerj.4654/fig-1)

Table 1 (A) Summary of sample information, genetic diversity indices and population specific F_{ST} for microsatellite loci of Australian smelt. (B) Summary of sample information for mitochondrial DNA sequences of Australian smelt. Number of samples used for genetic analysis (N), mean number of alleles per population (N_A), observed heterozygosity (H_O), expected heterozygosity (H_E), mean inbreeding index (F_{IS}).

(A)

Group name	Sampling site	Site code	Longitude (E)	Latitude (S)	N	N_A	H_O	H_E	F_{IS}	Population specific F_{ST}
SEQ-N	Tinana	MRD	152°42'57.8"	25°36'04.3"	26	8.90	0.686	0.733	0.066	0.196
SEQ-N	Mary_upper	MRU	152°48'47.9"	26°38'55.5"	29	10.90	0.779	0.826	0.059	0.136
SEQ-N	Noosa_lower	NSD	152°52'21.4"	26°17'05.7"	30	6.10	0.576	0.617	0.067	0.340
SEQ-N	Mooloolah_lower	MLD	153°0'44.64"	26°46'18.83"	16	4.60	0.670	0.604	-0.115	0.390
SEQ-N	Mooloolah_upper	MLU	152°55'13.1"	26°45'07.9"	32	6.40	0.609	0.566	-0.079	0.334
SEQ-S	Brisbane_lower	BRD	152°55'49.9"	27°30'16.05"	32	9.50	0.754	0.755	0.002	0.191
SEQ-S	Brisbane_upper	BRU	152°35'13.5"	27°58'43.9"	32	8.80	0.760	0.763	0.005	0.216
SEQ-S	Logan-Albert_lower	LAD	152°59'01.8"	28°10'15.6"	8	7.60	0.701	0.847	0.184	0.085
SEQ-S	Logan-Albert_upper	LAU	152°56'23.6"	28°19'19.7"	24	11.40	0.828	0.845	0.021	0.107
SEQ-S	Coomera_lower	CMD	153°11'20.9"	28°02'55.5"	24	13.90	0.839	0.887	0.054	0.074
SEQ-S	Coomera_upper	CMU	153°09'13.4"	28°05'01.8"	32	14.70	0.848	0.887	0.045	0.078
SEQ-S	Nerang_lower	NRD	153°17'52.0"	28°01'33.7"	8	6.40	0.718	0.798	0.106	0.156
SEQ-S	Nerang_upper	NRU	153°14'02.8"	28°07'29.2"	32	13.70	0.782	0.853	0.084	0.088
SEQ-S	Currumbin_lower	CRD	153°25'24.8"	28°10'41.9"	32	11.20	0.771	0.803	0.041	0.130
SEQ-S	Currumbin_upper	CRU	153°23'11.9"	28°12'49.6"	32	10.90	0.769	0.785	0.021	0.135

(B)

Group name	Sampling site	Site code	Longitude (E)	Latitude (S)	N (number of sample)
SEQ-N	Tinana	MRD	152°42'57.8"	25°36'04.3"	2
SEQ-N	Mary_upper	MRU	152°48'47.9"	26°38'55.5"	2
SEQ-N	Booloumba Creek	MBC	152°37'10.6"	26°41'02.5"	9
SEQ-N	Yabba Creek	MYC	152°38'39.5"	26°28'09.3"	8
SEQ-N	Noosa_lower	NSD	152°52'21.4"	26°17'05.7"	2
SEQ-N	Mooloolah_lower	MLD	153°0'44.64"	26°46'18.83"	2
SEQ-N	Mooloolah_upper	MLU	152°55'13.1"	26°45'07.9"	3
SEQ-S	Brisbane_lower	BRD	152°55'49.9"	27°30'16.05"	2
SEQ-S	Brisbane_upper	BRU	152°35'13.5"	27°58'43.9"	2
SEQ-S	Bundamba Creek	BDC	152°48'04.2"	27°36'03.9"	10
SEQ-S	Banks Creek	BSV	152°40'13.2"	27°26'36.9"	10
SEQ-S	Logan-Albert_lower	LAD	152°59'01.8"	28°10'15.6"	2
SEQ-S	Logan-Albert_upper	LAU	152°56'23.6"	28°19'19.7"	2
SEQ-S	Coomera_lower	CMD	153°11'20.9"	28°02'55.5"	2
SEQ-S	Coomera_upper	CMU	153°09'13.4"	28°05'01.8"	2
SEQ-S	Nerang_lower	NRD	153°17'52.0"	28°01'33.7"	2
SEQ-S	Nerang_upper	NRU	153°14'02.8"	28°07'29.2"	2
SEQ-S	Currumbin_lower	CRD	153°25'24.8"	28°10'41.9"	2
SEQ-S	Currumbin_upper	CRU	153°23'11.9"	28°12'49.6"	2

labelled PCR products were pooled and added to 10 μ l of Hi-Di™ formamide with 0.1 μ l of GeneScan™ 500 LIZ size standard. Fragment analysis was conducted on an ABI PRISM 3130 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA) according to the manufacturer's instructions. Data were scored using GENEMAPPER version 3.1 software (Applied Biosystems, Foster City, CA, USA).

Two individuals from each of the 15 populations represented in the microsatellite study were randomly selected for mtDNA analysis except the Mooloolah_upper (MLU) site from which one additional sample was included for analysis. Samples from four additional sites not included in microsatellite analysis were also sequenced—two sites from the Mary River and two sites from the Brisbane River (Fig. 1 and Table 1B). In total 68 individuals from nineteen sites were sequenced. A 666 bp fragment of the cytochrome *b* region of the mtDNA genome was selected for sequence analysis. The primers HYP5LA and HYP5HD (Thacker *et al.*, 2007) were used to amplify the region in 10 μ L reaction mixtures. PCR conditions were 4 min at 95 °C, followed by 45 cycles of 30 s at 95 °C, 45 s at 53 °C, 45 s at 72 °C and a final extension cycle of 7 min at 72 °C. MtDNA sequences were edited and aligned using Geneious version 9.1.5 (Kearse *et al.*, 2012).

Data analysis

Genetic diversity

Microsatellite genotype frequencies were checked for the presence of null alleles, large allele dropout and stuttering artefacts using MICRO-CHECKER v2.2.3 (Van Oosterhout *et al.*, 2004). Tests for linkage disequilibrium (LD) and departures of genotypic proportions expected under Hardy-Weinberg Equilibrium (HWE) were undertaken with exact tests for each population and over all loci using default settings in GENEPOP v4 (Rousset, 2008). Probability values were corrected using standard Bonferroni correction (Rice, 1989) whenever multiple testing was performed.

Genetic diversity averaged across ten loci within each of the 15 population samples was calculated from observed and expected heterozygosity using ARLEQUIN v3.5.1.2 (Excoffier & Lischer, 2010). Inbreeding index (F_{IS}) was estimated in FSTAT 2.9.3 (Goudet, 2001).

Population genetic structure

Genetic structure among the 15 populations was quantified through estimating pairwise and global F_{ST} values in ARLEQUIN. These were tested for significant deviation from panmictic expectations by 10,000 permutations of individuals among populations. Population-specific F_{ST} values were calculated using GESTE v2.0 (Foll & Gaggiotti, 2006) to evaluate the contribution of individual population samples to overall F_{ST} .

ARLEQUIN was used to evaluate the geographic structuring of genetic variation. F_{ST} was calculated for each locus separately and as a weighted average over the ten microsatellite loci. Statistical significance of F_{ST} was determined by 1,000 permutations of individuals among populations. Hierarchical structuring of variation was calculated using analyses of variance (AMOVA) in ARLEQUIN. Two hierarchical arrangements of the 15 populations were analysed where the highest level was either (a) two groups (species), SEQ-N (MRD, MRU, NSD, MLD, MLU) and SEQ-S (BRD, BRU, LAD, LAU, CMD, CMU, NRD, NRU, CRD, CRU) or (b) catchment division, grouped into eight rivers. These were: Mary (MRD,

MRU), Noosa (NSD), Mooloolah (MLD, MLU), Brisbane (BRD, BRU), Logan (LAD, LAU), Coomera (CMD, CMU), Nerang (NRD, NRU) and Currumbin (CMD, CMU). Three hierarchical levels of variation were analysed for each arrangement: among rivers (F_{CT}), among populations within rivers (F_{SC}) and among all populations (F_{ST}).

Bayesian clustering methods implemented in STRUCTURE v.2.3.1 (Pritchard, Stephens & Donnelly, 2000) were applied to estimate the number of genetically homogeneous clusters (Latch et al., 2006; Hasselman, Ricard & Bentzen, 2013). This programme builds genetic clusters by minimizing linkage disequilibrium and deviations from Hardy-Weinberg equilibrium expectations within clusters. All individuals were assigned to clusters without prior knowledge of their geographic origin using the admixture model with correlated allelic frequencies. Ten independent runs with the number of potential genetic clusters (K) from 1 to 16 were carried out to verify that the estimates of K were consistent across runs. The burn-in length was set at 250,000 iterations followed by a run phase of one million iterations. The generated results were imported into the software STRUCTURE HARVESTER (Earl & vonHoldt, 2012) to calculate the *ad hoc* ΔK statistic (Evanno, Regnaut & Goudet, 2005). The K value where ΔK had the highest value was identified as the most likely number of clusters.

Analysis of isolation by distance

A test for a positive association between genetic and geographic distances (Isolation by distance (IBD)) based on microsatellite DNA loci was carried out using a Mantel test (10,000 permutations) in ARLEQUIN. Genetic distance was represented as F_{ST} . Stream distances were calculated between river mouths and then sample sites using Google Earth.

Migration and gene flow

BAYESASS v1.3 was used to calculate contemporary migration rates over the past few generations, where m is the proportion of immigrants in a focal population i that arrive from a source population j (Wilson & Rannala, 2003). Migration rates were estimated for all pairs of sites and rate were reported that fell outside the 95% confidence interval simulated for uninformative data (Wilson & Rannala, 2003). We also used the Bayesian assignment procedure of Rannala & Mountain (1997), as implemented in GENECLASS 2 (Piry et al., 2004) to estimate whether our samples might contain individuals that were first generation (F_0) immigrants from unsampled populations. Here we used the Paetkau et al. (2004) method to compute probabilities from 10,000 simulated genotypes to identify F_0 immigrants.

Analysis of mtDNA sequence data

A neighbour-joining (NJ) tree analysis was performed using the HKY distance model in Geneious version 9.1.5 with 1,000 bootstrap replicates (Kearse et al., 2012). In addition to the 68 sequences generated from this study, two Genbank accessions were used, one representing *R. tasmanica*: JN232589; and one representing *R. semoni*: JN232588 (BurrIDGE et al., 2012). The *R. semoni* sequence JN232588 lacks locality information (C BurrIDGE, pers. comm., 2017), but likely belongs to a southern lineage of *R. semoni* which are known to have a closer mtDNA relationship with *R. tasmanica* than to northern lineages (Hughes et al., 2014).

RESULTS

Genetic variability and levels of differentiation

After Bonferroni correction, 3 out of 15 populations exhibited deviations from HWE in only one or two loci (Table S1). All loci were kept for further analyses since deviations were not consistent across populations. Instances of null alleles estimated using MICRO-CHECKER were rare and not consistently associated with specific loci or populations (Table S2). No evidence for genotypic linkage disequilibrium was observed between any pair of loci across all populations.

Population genetic diversity indices are shown in Table 1A. Microsatellite genetic diversity was high. Mean number of alleles per population ranged from 4.60 (MLD) to 14.70 (CMU) and heterozygosity averaged across loci ranged from 0.566 (MLU) to 0.887 (CMD and CMU). Most sites exhibited positive F_{IS} values, indicating that most of the populations had slight heterozygote deficits.

Most of the pairwise F_{ST} values between the 15 populations were significant and ranged from -0.018 to 0.404 . The SEQ-N populations were more diverged from one another than the populations in the SEQ-S group. The lowest pairwise F_{ST} value ($F_{ST} = -0.018$; $P < 0.05$) was observed between populations NRD and NRU. The highest genetic divergence ($F_{ST} = 0.404$; $P < 0.05$) was observed between populations NSD and MLU. Out of 105 comparisons, only two comparisons were non-significant ($P > 0.05$) and each of these pairs was from within the same river (Logan-Albert and Nerang). However, one site of each of these pairs contains a small number of samples which probably reduces the power to detect the significant F_{ST} values. Generally F_{ST} comparisons revealed much less divergence among populations within the same river than between populations from different rivers (Table 2).

STRUCTURE analysis revealed the highest likelihood was at $K = 8$ clusters (Average log probability of data $\ln[P(DK)] \pm SD = -15125.4 \pm 0.584618$) (Fig. 2, Table S3) indicating this as the best estimate of the true number of genetic clusters. The height of ΔK was used as an indicator of the strength of the signal detected by STRUCTURE (Evanno, Regnaut & Goudet, 2005). ΔK also showed the highest peak at $K = 8$, suggesting eight genetically homogeneous clusters across the sampled populations (Fig. S1) and negligible migration was observed among rivers (Fig. 2).

Strong genetic structure was observed by AMOVA. The AMOVA showed significant genetic differentiation between the two groups (SEQ-N and SEQ-S) ($F_{CT} = 0.05$), but also among populations within groups ($F_{SC} = 0.19$) (Table 3A). There were similar patterns observed in the two groups when each group (SEQ-N and SEQ-S) was analysed separately, with the F_{CT} value (among rivers) higher than the F_{SC} value (among sites within rivers) in both groups (Table 3Bi and 3Bii). However, the overall F_{ST} values, and each of the other F statistics in the hierarchy were slightly higher within the SEQ-N group than the SEQ-S group.

Isolation-by-distance

There was a significant correlation between genetic differentiation and stream distance among populations in the SEQ-S group ($R^2 = 0.3687$, $p = 0.001$; BRD, BRU, LAD, LAU,

Table 2 Pairwise F_{ST} values among all pairs of populations based on microsatellite data. F_{ST} estimates significance levels < 0.05 are in bold following Bonferroni correction. See Table 1A for site codes and locations.

	MRD	MRU	NSD	MLD	MLU	BRD	BRU	LAD	LAU	CMD	CMU	NRD	NRU	CRD	CRU
MRD	0.000														
MRU	0.108	0.000													
NSD	0.358	0.310	0.000												
MLD	0.250	0.182	0.362	0.000											
MLU	0.318	0.244	0.404	0.009	0.000										
BRD	0.178	0.131	0.323	0.203	0.276	0.000									
BRU	0.173	0.136	0.320	0.218	0.295	0.013	0.000								
LAD	0.243	0.101	0.394	0.248	0.316	0.157	0.170	0.000							
LAU	0.244	0.137	0.368	0.246	0.317	0.173	0.189	-0.006	0.000						
CMD	0.212	0.151	0.319	0.186	0.259	0.130	0.146	0.126	0.161	0.000					
CMU	0.203	0.141	0.293	0.165	0.236	0.124	0.139	0.127	0.164	0.010	0.000				
NRD	0.261	0.125	0.393	0.244	0.316	0.187	0.191	0.073	0.082	0.152	0.141	0.000			
NRU	0.229	0.110	0.329	0.203	0.261	0.176	0.183	0.053	0.074	0.146	0.137	-0.018	0.000		
CRD	0.263	0.152	0.353	0.222	0.277	0.195	0.211	0.125	0.147	0.166	0.156	0.076	0.062	0.000	
CRU	0.305	0.178	0.397	0.270	0.320	0.248	0.256	0.120	0.150	0.215	0.201	0.073	0.059	0.011	0.000




Figure 2 Results from structure indicating individual assignment and population clustering of Australian smelt for 389 individuals from all 15 locations sampled (Table 1A) in south-east Queensland, Australia. Individual sampling locations is listed below the figure and see Table 1A for site codes and locations. Each column represents one individual and the colours represent the probability membership coefficient of that individual for each genetic cluster. Results of Evanno's Mean LnP(D) and ΔK (Table S3; Fig. S1) indicate that the best supported K values for all populations are 8.

Full-size DOI: 10.7717/peerj.4654/fig-2

Table 3 Analyses of molecular variance (AMOVA) for hierarchical arrangements of the 15 sampling sites based on microsatellite data. The hierarchical levels tested were among rivers (F_{CT}), among populations within rivers (F_{SC}) and among all populations (F_{ST}). The total genetic variation is shown as a percentage for each hierarchical partitioning. Hierarchical arrangement of sites: (a) sites divided into two groups: SEQ-N & SEQ-S; (b) sites divided into catchment divisions within group.

Structure tested		Observed partition		F-Statistics
		Variance	% of variation	
(a)	Based on group (SEQ-N & SEQ-S)			
	Between groups	0.07255 Va	5.45	$F_{CT} = 0.05^*$
	Among sites within group	0.23313 Vb	17.50	$F_{SC} = 0.19^*$
	Within sites	1.02632 Vc	77.05	$F_{ST} = 0.23^*$
(b)	Based on river			
(i)	SEQ-N group			
	Among rivers	0.17274 Va	19.23	$F_{CT} = 0.19^*$
	Among sites within rivers	0.06542 Vb	7.28	$F_{SC} = 0.09^*$
	Within sites	0.66026 Vc	73.49	$F_{ST} = 0.27^*$
(ii)	SEQ-S group			
	Among rivers	0.27139 Va	12.49	$F_{CT} = 0.13^*$
	Among sites within rivers	0.01106 Vb	0.51	$F_{SC} = 0.006^*$
	Within sites	1.89107 Vc	87.01	$F_{ST} = 0.13^*$

Notes.

* $P < 0.001$.

CMD, CMU, NRD, NRU, CRD, and CRU) (Fig. 3A), but not for the SEQ-N group ($R^2 = 0.0355$, $p = 0.302$; MRD, NSD, MRU, MLD and MLU) (Fig. 3B).

Contemporary migration

Very little contemporary migration was observed among the coastal river populations. Only six sampled populations contained individuals that were identified as potential immigrants from the BAYESASS analysis. In all cases, the putative source population was the paired site within the same catchment (Table 4). In most cases, dispersal was from the upstream to the downstream site. Only individuals from Currumbin Creek were estimated to have




Figure 3 (A) Analysis of isolation by distance for SEQ-S populations. (B) Analysis of isolation by distance for SEQ-N populations.

Full-size DOI: [10.7717/peerj.4654/fig-3](https://doi.org/10.7717/peerj.4654/fig-3)

dispersed in an upstream direction. Only thirty-six (<10%) of 389 individuals across all sites were identified as F_0 migrants using the “detection of first generation migrants” option in GENECLASS2 (Table 5).

MtDNA sequences analysis

The edited alignment for the *cyt b* gene was 575 bp and included 121 variable positions. All sequences are lodged under GenBank accession numbers [MG867590–MG867657](#). The neighbour-joining tree revealed two strongly supported clades (bootstrap 89% SEQ-S; 96% SEQ-N; Fig. 4). Phylogeographic structure was also clearly evident within clades. All individuals from four sites in the Brisbane River formed a distinct clade, and all three rivers sampled for the SEQ-N lineage formed shallow clades (i.e., Mary, Noosa and Mooloolah rivers; Fig. 4). Genetic distance was high between northern smelt lineages and the southern smelt sequences used as outgroups (uncorrected mean nucleotide distance 0.15–0.17). The mean nucleotide distance between the two northern lineages (SEQ-N and SEQ-S) was 0.04 (SE = 0.007).

DISCUSSION

Population structure and dispersal

Based on previous studies of Australian smelt ([Woods et al., 2010](#); [Hughes et al., 2014](#)), we had hypothesized that *R. semoni* in south-east Queensland would exhibit limited genetic connectivity among river systems. Our findings of strong genetic differentiation among rivers support this hypothesis. For both groups (SEQ-N and SEQ-S), there were highly significant F_{ST} values, which indicated that populations were not panmictic. The results of the significant genetic differentiation among rivers were also consistent with the results of population structure revealed from the STRUCTURE analysis, suggesting restricted gene flow and limited dispersal among populations in both groups. Limited dispersal

Table 4 Estimates of contemporary migration rates (m) for each populations based on microsatellite data. Migration rate estimates were based on unidirectional assignment of microsatellite genotypes in BAYESASS v1.3. Proportion of nonmigrant values represents the proportion of individuals assigned back to their sampling site of origin. Mean and 95% CI (confidence interval) values for uninformative data were 0.833 (0.675, 0.992). Putative source of migrants represents source population supplying migrants into the focal population where the estimated migration rate exceeds the upper 95% CI value for uninformative data (0.0124). Migration rates and source of migrants were not provided for cases where the 95% CI of estimated migration rate overlapped with the 95% CI value for uninformative data.

Focal populations (Site code)	Proportion of nonmigrants	Putative source of migrants	Migration rate (m), mean (95% CI)
Tinana Creek (MRD)	0.988	–	–
Mary_upper (MRU)	0.987	–	–
Noosa_lower (NSD)	0.990	–	–
Mooloolah_lower (MLD)	0.685	Mooloolah_upper (MLU)	0.240 (0.130, 0.317)
Mooloolah_upper (MLU)	0.990	–	–
Brisbane_lower (BRD)	0.677	Brisbane_upper (BRU)	0.287 (0.232, 0.325)
Brisbane_upper (BRU)	0.989	–	–
Logan/Albert_lower (LAD)	0.699	Logan/Albert_upper (LAU)	0.145 (0.033, 0.279)
Logan/Albert_upper (LAU)	0.987	–	–
Coomera_lower (CMD)	0.681	Coomera_upper (CMU)	0.271 (0.198, 0.323)
Coomera_upper (CMU)	0.990	–	–
Nerang_lower (NRD)	0.668	Nerang_upper (NRU)	0.148 (0.036, 0.281)
Nerang_upper (NRU)	0.990	–	–
Currumbin_lower (CRD)	0.990	–	–
Currumbin_upper (CRU)	0.677	Currumbin_lower (CRD)	0.286 (0.235, 0.326)

was supported by our first-generation migrant detection analysis in GeneClass2, which demonstrated that less than 10% of individuals in each population were immigrants.

The sample from Tinana Creek (MRD site), was differentiated from the rest of the populations in the SEQ-N group (Table 2). This might be the result of a barrier which separates Tinana Creek from the rest of the Mary River system despite their close proximity to one another. Tinana Creek runs into the Mary River not far from the mouth, with both drainages having tidal estuarine reaches in the lower sections. The differentiation of the Tinana Creek population from the main stem of the Mary River is also observed in a number of other freshwater species including Mary River Cod, *Maccullochella mariensis* (Huey, Espinoza & Hughes, 2013), Mary River Turtle, *Elusor macrurus* (Schmidt et al., 2018), freshwater crayfish *Cherax dispar* (Bentley, Schmidt & Hughes, 2010) and Australian lungfish *Neoceratodus fosteri* (Hughes et al., 2015).

Although populations in the SEQ-N group were slightly more highly structured than those in the SEQ-S group, fishes in both groups exhibited restricted gene flow. The high genetic structuring in SEQ-N populations might be the result of genetic divergence within the lineage that occurs over a very small geographical range. The lower levels of genetic structure of *R. semoni* populations in the SEQ-S group than the SEQ-N group suggests that more current or recent gene flow occurs within and among the catchments in SEQ-S group than for the SEQ-N group. This could be because the catchments in the SEQ-S group have

Table 5 Results of the assessment for detecting first-generation migrants performed using GENECLASS2 showing the number of individual migrants ($P < 0.05$) detected per sampling location. Results are based on the $L_{\text{home}}/L_{\text{max}}$ statistic for microsatellite data. $L_{\text{home}}/L_{\text{max}}$ is the ratio of the likelihood computed from the population where the individual was sampled (L_{home}) over the highest likelihood value among all population samples including the population where the individual was sampled. F_0 is the first generation migrant. See Table 1A for site codes and locations.

Sample to	F_0 Migrants from															
	MRD	MRU	NSD	MLD	MLU	BRD	BRU	LAD	LAU	CMD	CMU	NRD	NRU	CRD	CRU	
MRD	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
MRU	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
NSD	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
MLD	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	
MLU	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
BRD	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	
BRU	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
LAD	0	0	0	0	0	0	0	4	0	0	0	0	0	0	0	
LAU	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	
CMD	0	0	0	0	0	0	0	0	0	6	0	0	0	0	0	
CMU	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
NRD	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	
NRU	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	
CRD	0	0	0	0	0	0	0	0	0	0	0	0	0	6	0	
CRU	0	0	0	0	0	0	0	0	0	0	0	0	0	4	0	




Figure 4 Neighbour-joining tree of the *cyt b* dataset for 68 Australian smelt samples from 19 sampling localities. Individual sample codes coloured according to river. Node values are bootstrap support. See Table 1B for site codes and locations. Individual sample codes coloured according to river. Node values are bootstrap support. See Table 1B for site codes and locations.

Full-size DOI: 10.7717/peerj.4654/fig-4

been connected more recently and/or more often than those in the SEQ-N group. This could be because during extreme flood events, SEQ-S catchments might be connected by freshwater plumes flowing into Southern Moreton Bay. Another plausible reason might be that these cryptic species (SEQ-N and SEQ-S) have different microhabitat preferences, with SEQ-N being less tolerant of brackish conditions (Hughes, 2015).

An important model for stream dwelling species is isolation by distance (IBD). In this model, equilibrium between genetic drift and gene flow may be reached in species where the life time dispersal distance is less than the range. Here, a relationship between stream

distance and genetic differentiation should be evident (*Wright, 1943*). In this study, a strong IBD relationship was identified among the SEQ-S populations, but not among SEQ-N populations. This suggests that for SEQ-S populations, dispersal, when it occurs, is more likely between nearby catchments. Similar IBD relationships have been reported for other coastline restricted species (*Keenan, 1994; Jerry & Baverstock, 1998; Shaddick et al., 2011; Schmidt et al., 2014*). Lack of IBD for the SEQ-N group may be attributed to an insufficient number of population samples available for comparison and/or the greater degree of population isolation within this group relative to the SEQ-S group, consistent with the overall higher F_{ST} estimates among SEQ-N populations. *Hughes et al. (2014)* observed similarly contrasting patterns of population genetic structure between cryptic species groups of southern Australian smelt. In that study, two informal species groups (MTV and SEC) with adjacent distributions along the western and eastern coast of southern Victoria had microsatellite-based F_{ST} values of 0.19 and 0.07 respectively (*Hughes et al., 2014*). Using otolith microchemistry, *Hughes et al. (2014)* also showed that the more structured western group (MTV) had a greater proportion of nondiadromous populations relative to the weaker structured eastern group (SEC). The similar pattern of contrasting structure observed here between northern groups in the Australian smelt complex (SEQ-N and SEQ-S), is probably not due to differences in diadromous behaviour because preliminary evidence from otolith chemistry suggests all of these populations are nondiadromous (*R Islam, 2017, unpublished data*). Higher structuring of the SEQ-N group could possibly be due to genetic drift if these populations have been established for a longer period of time at the northern-most limit of Australian smelt distribution relative to the SEQ-S populations.

The complementary pattern of population structuring in both microsatellite and mtDNA data between the SEQ-N and SEQ-S groups could reflect long-term phylogeographic isolation or perhaps further evidence for cryptic species within Australian smelts as first highlighted by *Hammer et al. (2007)*. Mean *cyt b* divergence of 4% between SEQ-N and SEQ-S samples is close to the 3.6% divergence observed for the full mitochondrial molecule reported by *Schmidt, Islam & Hughes (2016)*, and is within the range of lineage divergence reported for *R. semoni* in southern Queensland (*Page & Hughes, 2014*). It should be noted that *Schmidt, Islam & Hughes (2016)* incorrectly attributed mitochondrial sequence data for the SEQ-N group to the CEQ group (i.e., Genbank accession [KX421785](#) corresponds to SEQ-N and [KX421784](#) corresponds to SEQ-S). The level of *cyt b* divergence between the SEQ groups and southern lineages of *R. semoni* is very large (15–17%) and adds to previous studies that have highlighted the likely existence of a cryptic species complex within the taxon currently referred to as *R. semoni* (*Hammer et al., 2007; Hughes et al., 2014*).

Contemporary migration

The Bayesian assignment analysis detected contemporary movement of individuals only between proximate sites within rivers ([Table 4](#)). Contemporary dispersal was not observed between rivers. Although, most of the sites that we sampled were within 10–60 km stream distance of another sampled site, there was no contemporary dispersal among the majority of those rivers in either group. The present results coincide with the findings of some southern smelt populations, where contemporary movement was observed among

populations within a catchment (*Woods et al., 2010*). The species appears to occur in a wide range of freshwater habitats, many of which are isolated by long stretches of unfavourable habitat. Our data therefore suggest that if local extinctions occur in one or more of these streams within a reach of the river, then recolonization from elsewhere is unlikely to occur rapidly.

CONCLUSION

Finally, the present study revealed high levels of population structuring within and between drainages, which suggested that contemporary movement among sites was rare and limited to sites within the same river. Little conservation attention has been given to the Australian smelt since it has long been considered a common species distributed widely across south-eastern Australia. The findings of the present study and other recent research (*Hammer et al., 2007; Crook, Macdonald & Raadik, 2008; Hughes et al., 2014*) suggest that southern smelts are a genetically complex and ecologically diverse taxonomic group. Therefore, proper conservation and management will require appropriate taxonomic treatment to align species with the clear genetic divisions now recognised across the range of Australian smelt.

Further, eight isolated management units (MUs) along the south-east Queensland drainages were detected in *R. semoni* from the microsatellite dataset (*Fig. 2*) demonstrating little to no gene flow between them. These management units align with individual coastal catchments, which suggest that other genetically distinct populations may exist in coastal rivers not sampled in this study. High levels of genetic divergence between the two lineages have important implications for the conservation of these endemic freshwater cryptic fish species. Therefore, findings of the present study on population structure of Australian smelt will help to formulate effective management and conservation plans for this cryptic species complex across their geographic range.

ACKNOWLEDGEMENTS

We thank Leo Lee, Nathan McIntyre, Dale Bryan Brown, Juan Tao for help in collecting samples. Thanks to Kathryn Real and Jemma Somerville for assistance with genetic laboratory work.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Funding for this research was provided by Griffith University to Md Rakeb-Ul Islam through International Postgraduate Research Award. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Griffith University.

Competing Interests

Jane M. Hughes is an Academic Editor of PeerJ.

Author Contributions

- Md Rakeb-Ul Islam conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, approved the final draft.
- Daniel J. Schmidt analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- David A. Crook authored or reviewed drafts of the paper, approved the final draft.
- Jane M. Hughes conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All field and experimental protocols carried out in this study were approved by the Griffith University Animal Ethics Committee.

Data Availability

The following information was supplied regarding data availability:

Microsatellite raw data and GenBank accession numbers of Cyt b haplotypes are available in [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4654#supplemental-information>.

REFERENCES

- Amoros C, Bornette G. 2002.** Connectivity and biocomplexity in waterbodies of riverine floodplains. *Freshwater Biology* **47**:761–776 DOI [10.1046/j.1365-2427.2002.00905.x](https://doi.org/10.1046/j.1365-2427.2002.00905.x).
- Baumgartner LJ, Harris JH. 2007.** Passage of non-salmonid fish through a Deelder lock on a lowland river. *River Research and Applications* **23**(10):1058–1069 DOI [10.1002/rra.1032](https://doi.org/10.1002/rra.1032).
- Bentley AI, Schmidt DJ, Hughes JM. 2010.** Extensive intraspecific genetic diversity of a freshwater crayfish in a biodiversity hotspot. *Freshwater Biology* **55**:1861–1873 DOI [10.1111/j.1365-2427.2010.02420.x](https://doi.org/10.1111/j.1365-2427.2010.02420.x).
- Bernays SJ, Schmidt DJ, Hurwood DA, Hughes JM. 2015.** Phylogeography of two freshwater prawn species from far-northern Queensland. *Marine and Freshwater Research* **66**:256–266 DOI [10.1071/MF14124](https://doi.org/10.1071/MF14124).
- Burridge CP, Craw D, Jack DC, King TM, Waters JM. 2008.** Does fish ecology predict dispersal across a river drainage divide? *Evolution* **62**:1484–1499 DOI [10.1111/j.1558-5646.2008.00377.x](https://doi.org/10.1111/j.1558-5646.2008.00377.x).

- Burridge CP, McDowall RM, Craw D, Wilson MVH, Waters JM. 2012.** Marine dispersal as a pre-requisite for Gondwanan vicariance among elements of the galaxiid fish fauna. *Journal of Biogeography* **39**:306–321 DOI [10.1111/j.1365-2699.2011.02600.x](https://doi.org/10.1111/j.1365-2699.2011.02600.x).
- Crook DA, Macdonald JJ, Raadik TA. 2008.** Evidence of diadromous movements in a coastal population of southern smelts (Retropinninae: *Retropinna*) from Victoria, Australia. *Marine and Freshwater Research* **59**:638–646 DOI [10.1071/MF07238](https://doi.org/10.1071/MF07238).
- Earl DA, vonHoldt BM. 2012.** STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetic Resources* **4**:359–361 DOI [10.1007/s12686-011-9548-7](https://doi.org/10.1007/s12686-011-9548-7).
- Evanno G, Regnaut S, Goudet J. 2005.** Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* **14**:2611–2620 DOI [10.1111/j.1365-294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x).
- Excoffier L, Lischer HEL. 2010.** Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* **10**(3):564–567 DOI [10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x).
- Foll M, Gaggiotti O. 2006.** Identifying the environmental factors that determine the genetic structure of populations. *Genetics* **174**:875–891 DOI [10.1534/genetics.106.059451](https://doi.org/10.1534/genetics.106.059451).
- Geist J. 2011.** Integrative freshwater ecology and biodiversity conservation. *Ecological Indicators* **11**:1507–1516 DOI [10.1016/j.ecolind.2011.04.002](https://doi.org/10.1016/j.ecolind.2011.04.002).
- Goudet J. 2001.** FSTAT, version 2.9.3, A program to estimate and test gene diversities and fixation indices. Lausanne: Lausanne University.
- Hammer MP, Adams M, Unmack PJ, Walker KF. 2007.** A rethink on *Retropinna*: conservation implications of new taxa and significant genetic sub-structure in Australian smelts (Pisces: Retropinnidae). *Marine and Freshwater Research* **58**:327–341 DOI [10.1071/MF05258](https://doi.org/10.1071/MF05258).
- Hasselman DJ, Ricard D, Bentzen P. 2013.** Genetic diversity and differentiation in a wide ranging anadromous fish, American shad (*Alosa sapidissima*), is correlated with latitude. *Molecular Ecology* **22**:1558–1573 DOI [10.1111/mec.12197](https://doi.org/10.1111/mec.12197).
- Huey JA, Baker AM, Hughes JM. 2010.** High levels of genetic structure in the Australian freshwater fish, *Ambassis macleayi*. *Journal of the North American Benthological Society* **29**(3):1148–1160 DOI [10.1899/09-093.1](https://doi.org/10.1899/09-093.1).
- Huey JA, Espinoza T, Hughes JM. 2013.** Natural and anthropogenic drivers of genetic structure and low genetic variation in the endangered freshwater cod, *Maccullochella mariensis*. *Conservation Genetics* **14**:997–1008 DOI [10.1007/s10592-013-0490-y](https://doi.org/10.1007/s10592-013-0490-y).
- Hughes JM. 2007.** Constraints on recovery: using molecular methods to study connectivity of aquatic biota in rivers and streams. *Freshwater Biology* **52**(4):616–631 DOI [10.1111/j.1365-2427.2006.01722.x](https://doi.org/10.1111/j.1365-2427.2006.01722.x).
- Hughes JM. 2015.** Current and future challenges of conserving freshwater biodiversity: a molecular perspective. *Pertanika Journal of Tropical Agricultural Science* **38**(4):413–426.

- Hughes JM, Schmidt DJ, Finn DS. 2009.** Genes in streams: using DNA to understand the movement of freshwater fauna and their riverine habitat. *BioScience* **59**(7):573–583 DOI [10.1525/bio.2009.59.7.8](https://doi.org/10.1525/bio.2009.59.7.8).
- Hughes JM, Schmidt DJ, Huey JA, Real KM, Espinoza T, McDougall A, Kind PK, Brooks S, Roberts DT. 2015.** Extremely low microsatellite diversity but distinct population structure in a long-lived threatened species, the Australian lungfish *Neoceratodus fosteri* (Dipnoi). *PLOS ONE* **10**(4):e0121858 DOI [10.1371/journal.pone.0121858](https://doi.org/10.1371/journal.pone.0121858).
- Hughes JM, Schmidt DJ, Macdonald JI, Huey JA, Crook DA. 2014.** Low interbasin connectivity in a facultatively diadromous fish: evidence from genetics and otolith chemistry. *Molecular Ecology* **23**:1000–1013 DOI [10.1111/mec.12661](https://doi.org/10.1111/mec.12661).
- Islam MR-U, Schmidt DJ, Hughes JM. 2017.** Development and characterization of 21 novel microsatellite markers for the Australian smelt *Retropinna semoni* (Weber, 1895). *Journal of Applied Ichthyology* **33**:824–828 DOI [10.1111/jai.13391](https://doi.org/10.1111/jai.13391).
- Jerry DR, Baverstock PR. 1998.** Consequences of a catadromous life-strategy for levels of mitochondrial DNA differentiation among populations of the Australian bass, *Macquaria novemaculeata*. *Molecular Ecology* **7**:1003–1013 DOI [10.1046/j.1365-294x.1998.00418.x](https://doi.org/10.1046/j.1365-294x.1998.00418.x).
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Mentjies P, Drummond A. 2012.** Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**(12):1647–1649 DOI [10.1093/bioinformatics/bts199](https://doi.org/10.1093/bioinformatics/bts199).
- Keenan CP. 1994.** Recent evolution of population structure in Australian barramundi, *Lates calcarifer* (Bloch): an example of isolation by distance in one dimension. *Australian Journal of Marine and Freshwater Research* **45**:1123–1148 DOI [10.1071/MF9941123](https://doi.org/10.1071/MF9941123).
- Latch EK, Dharmarajan G, Glaubitz JC, Rhodes OE. 2006.** Relative performance of Bayesian clustering software for inferring population substructure and individual assignment at low levels of population differentiation. *Conservation Genetics* **7**:295–302 DOI [10.1007/s10592-005-9098-1](https://doi.org/10.1007/s10592-005-9098-1).
- Leclerc E, Mailhot Y, Mingelbier M, Bernatchez L. 2008.** The landscape genetics of yellow perch (*Perca flavescens*) in a large fluvial ecosystem. *Molecular Ecology* **17**:1702–1717 DOI [10.1111/j.1365-294X.2008.03710.x](https://doi.org/10.1111/j.1365-294X.2008.03710.x).
- McDowall RM. 1996.** Family Retropinnidae: southern smelts. In: McDowall RM, ed. *Freshwater fishes of South-eastern Australia*. Chatswood: Reed books, 92–95.
- McGlashan DJ, Hughes JM. 2000.** Reconciling patterns of genetic variation with stream structure, earth history and biology in the Australian freshwater fish *Craterocephalus stercusmuscarum* (Athernidae). *Molecular Ecology* **9**:1737–1751 DOI [10.1046/j.1365-294x.2000.01054.x](https://doi.org/10.1046/j.1365-294x.2000.01054.x).
- McGlashan DJ, Hughes JM. 2002.** Extensive genetic divergence among populations of the Australian freshwater fish, *Pseudomugil signifer* (Pseudomugilidae),

- at different hierarchical scales. *Marine and Freshwater Research* **53**:897–907 DOI [10.1071/MF01107](https://doi.org/10.1071/MF01107).
- Paetkau D, Slade R, Burden M, Estoup A. 2004.** Genetic assignment methods for the direct, real-time estimation of migration rate: a simulation-based exploration of accuracy and power. *Molecular Ecology* **13**:55–65 DOI [10.1046/j.1365-294X.2004.02008.x](https://doi.org/10.1046/j.1365-294X.2004.02008.x).
- Page TJ, Hughes JM. 2014.** Contrasting insights provided by single and multispecies data in a regional comparative phylogeographic study. *Biological Journal of the Linnean Society* **111**:554–569 DOI [10.1111/bij.12231](https://doi.org/10.1111/bij.12231).
- Pereira LHG, Foresti F, Oliveira C. 2009.** Genetic structure of the migratory catfish *Pseudoplatystoma corruscans* (Siluriformes: Pimelodidae) suggesting homing behaviour. *Ecology of Freshwater Fish* **18**:215–225 DOI [10.1111/j.1600-0633.2008.00338.x](https://doi.org/10.1111/j.1600-0633.2008.00338.x).
- Piry S, Alapetite A, Cornuet J-M, Peatkau D, Baudouin L, Estoup A. 2004.** GENECLASS2: a software for genetic assignment and first-generation migrant detection. *Journal of Heredity* **95**:536–539 DOI [10.1093/jhered/esh074](https://doi.org/10.1093/jhered/esh074).
- Pritchard JK, Stephens M, Donnelly P. 2000.** Inference of population structure using multilocus genotype data. *Genetics* **155**:945–959.
- Pusey BJ, Kennard M, Arthington A. 2004.** *Freshwater fishes of north-eastern Australia*. Collingwood: CSIRO Publishing.
- Rannala B, Mountain JL. 1997.** Detecting immigration by using multilocus genotypes. *Proceedings of the National Academy of Sciences of the United States of America* **94**:9197–9201 DOI [10.1073/pnas.94.17.9197](https://doi.org/10.1073/pnas.94.17.9197).
- Rice WR. 1989.** Analyzing tables of statistical tests. *Evolution* **43**:223–225 DOI [10.2307/2409177](https://doi.org/10.2307/2409177).
- Rolls RJ. 2011.** The role of life-history and location of barriers to migration in the spatial distribution and conservation of fish assemblages in a coastal river system. *Biological Conservation* **144**(1):339–349 DOI [10.1016/j.biocon.2010.09.011](https://doi.org/10.1016/j.biocon.2010.09.011).
- Rousset F. 2008.** Genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources* **8**(1):103–106 DOI [10.1111/j.1471-8286.2007.01931.x](https://doi.org/10.1111/j.1471-8286.2007.01931.x).
- Schmidt DJ, Crook DA, Macdonald JI, Huey JA, Zampatti BP, Chilcott S, Raadik TA, Hughes JM. 2014.** Migration history and stock structure of two putatively diadromous teleost fishes as determined by genetic and otolith chemistry analyses. *Freshwater Science* **33**(1):193–206 DOI [10.1086/674796](https://doi.org/10.1086/674796).
- Schmidt DJ, Espinoza T, Connell M, Hughes JM. 2018.** Conservation genetics of the Mary River turtle (*Elusor macrurus*) in natural and captive populations. *Aquatic Conservation: Marine and Freshwater Ecosystems* **28**(1):115–123 DOI [10.1002/aqc.2851](https://doi.org/10.1002/aqc.2851).
- Schmidt DJ, Islam MR-U, Hughes JM. 2016.** Complete mitogenomes for two lineages of the Australian smelt, *Retropinna semoni* (Osmeriformes: Retropinnidae). *Mitochondrial DNA Part B* **1**(1):615–616 DOI [10.1080/23802359.2016.1209097](https://doi.org/10.1080/23802359.2016.1209097).
- Shaddick K, Gilligan DM, Burrridge CP, Jerry DR, Truong K, Beheregaray LB. 2011.** Historic divergence with contemporary connectivity in a catadromous fish, the estuary perch (*Macquaria colonorum*). *Canadian Journal of Fisheries and Aquatic Sciences* **68**:304–318 DOI [10.1139/F10-139](https://doi.org/10.1139/F10-139).

- Sharma S, Hughes JM. 2009.** Genetic structure and phylogeography of freshwater shrimps (*Macrobrachium australiense* and *Macrobrachium tolmerum*): the role of contemporary and historical events. *Marine and Freshwater Research* **60**:541–553 DOI [10.1071/MF07235](https://doi.org/10.1071/MF07235).
- Thacker CE, Unmack PJ, Matsui L, Rifenburg N. 2007.** Comparative phylogeography of five sympatric *Hypseleotris* species (Teleostei: Eleotridae) in southeastern Australia reveals a complex pattern of drainage basin exchanges with little congruence across species. *Journal of Biogeography* **34**(9):1518–1533 DOI [10.1111/j.1365-2699.2007.01711.x](https://doi.org/10.1111/j.1365-2699.2007.01711.x).
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004.** MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**(3):535–538 DOI [10.1111/j.1471-8286.2004.00684.x](https://doi.org/10.1111/j.1471-8286.2004.00684.x).
- Ward RD, Woodward M, Skibinski DOF. 1994.** A comparison of genetic diversity levels in marine, freshwater and anadromous fishes. *Journal of Fish Biology* **44**:213–232 DOI [10.1111/j.1095-8649.1994.tb01200.x](https://doi.org/10.1111/j.1095-8649.1994.tb01200.x).
- Wilson GA, Rannala B. 2003.** Bayesian inference of recent migration rates using multilocus genotypes. *Genetics* **163**:1177–1191.
- Woods RJ, Macdonald JI, Crook DA, Schmidt DJ, Hughes JM. 2010.** Contemporary and historical patterns of connectivity among populations of an inland river fish species inferred from genetics and otolith chemistry. *Canadian Journal of Fisheries and Aquatic Sciences* **67**:1098–1115 DOI [10.1139/F10-043](https://doi.org/10.1139/F10-043).
- Wright S. 1943.** Isolation by distance. *Genetics* **28**(2):114–138.