



Charles Darwin University

4-(Prop-2-yn-1-yloxy)benzene-1,2-dicarbonitrile

Chin, Yee Jan; Tan, Ai Ling; Wimmer, Franz L.; Mirza, Aminul Huq; Young, David J.; Ng, Seik Weng; Tiekink, Edward R T

Published in:
Acta Crystallographica Section E: Structure Reports Online

DOI:
[10.1107/S1600536812028309](https://doi.org/10.1107/S1600536812028309)

Published: 01/07/2012

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Chin, Y. J., Tan, A. L., Wimmer, F. L., Mirza, A. H., Young, D. J., Ng, S. W., & Tiekink, E. R. T. (2012). 4-(Prop-2-yn-1-yloxy)benzene-1,2-dicarbonitrile. *Acta Crystallographica Section E: Structure Reports Online*, E68(Part 7), o2293-o2294. <https://doi.org/10.1107/S1600536812028309>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-(Prop-2-yn-1-yloxy)benzene-1,2-dicarbonitrile

 Yee Jan Chin,^a Ai Ling Tan,^a Franz L. Wimmer,^a
 Aminul Huq Mirza,^a David J. Young,^{a,‡} Seik Weng Ng^{b,c}
 and Edward R. T. Tiekink^{b,*}

^aFaculty of Science, Universiti Brunei Darussalam, Jalan Tungku Link BE 1410, Negara, Brunei Darussalam, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department and Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia
 Correspondence e-mail: edward.tiekink@gmail.com


Received 19 June 2012; accepted 22 June 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.059; wR factor = 0.163; data-to-parameter ratio = 13.0.

In the title compound, $\text{C}_{11}\text{H}_6\text{N}_2\text{O}$, the complete molecule is generated by the application of crystallographic twofold symmetry (the molecule is disordered about this axis). The prop-2-yn-1-yl residue is slightly twisted out of the plane of the benzene ring [$\text{C}-\text{O}-\text{C}-\text{C}$ torsion angle = $173.1(3)^\circ$] and is orientated away from the nitrile substituents. In the crystal, supramolecular chains along the a axis, arising from $\text{C}-\text{H}\cdots\text{N}$ interactions, are connected into stacks along the c axis by $\pi-\pi$ interactions between the benzene rings [centroid-centroid distance = $3.6978(6)$ Å = length of the c axis].

Related literature

For the solubilization and some applications of phthancyanine dyes, see: Jiang *et al.* (2011); Slevin *et al.* (2001). For the synthesis of substituted phthalonitriles, see: Wöhrle *et al.* (1993); Wu *et al.* (1998); Li & Lieberman (2001); Slevin *et al.* (2001); Li *et al.* (2008); Seven *et al.* (2009); Foo *et al.* (2012).



‡ Additional correspondence author: david.young@ubd.edu.bn.

Experimental

Crystal data

$\text{C}_{11}\text{H}_6\text{N}_2\text{O}$
 $M_r = 182.18$
 Monoclinic, $C2/m$
 $a = 11.4809(9)$ Å
 $b = 22.2091(16)$ Å
 $c = 3.6978(6)$ Å
 $\beta = 91.304(10)^\circ$

$V = 942.62(18)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100$ K
 $0.15 \times 0.05 \times 0.05$ mm

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
 Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2012)
 $T_{\min} = 0.792$, $T_{\max} = 1.000$

3258 measured reflections
 1114 independent reflections
 840 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.163$
 $S = 1.08$
 1114 reflections
 86 parameters
 12 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.41$ e Å⁻³
 $\Delta\rho_{\min} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C3}-\text{H3}\cdots\text{N1}^i$	0.95	2.62	3.509 (3)	155

Symmetry code: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors gratefully acknowledge funding from the Brunei Research Council, and thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6862).

References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
 Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Foo, C. C., Tan, A. L., Wimmer, F. L., Mirza, A. H., Young, D. J., Ng, S. W. & Tiekink, E. R. T. (2012). *Acta Cryst.* **E68**, o601.
 Jiang, X.-J., Yeung, S.-L., Lo, P.-C., Fong, W. P. & Ng, D. K. P. (2011). *J. Med. Chem.* **54**, 320–330.
 Li, H., Jensen, T. J., Fronczek, F. R. & Vicente, G. H. (2008). *J. Med. Chem.* **51**, 502–511.
 Li, Z. & Lieberman, M. (2001). *Inorg. Chem.* **40**, 932–939.
 Seven, O., Dindar, B. & Gultekin, B. (2009). *Turk. J. Chem.* **33**, 123–134.

- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sleven, J., Görller-Walrand, C. & Binnemans, K. (2001). *Mater. Sci. Eng. C*, **18**, 229–238.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Wöhrle, D., Eskes, M., Shigehara, K. & Yamada, A. (1993). *Synthesis*, pp. 194–196.
- Wu, Y., Tian, H., Chen, K., Liu, Y. & Zhu, D. (1998). *Dyes Pigm.* **37**, 317–325.

supporting information

Acta Cryst. (2012). E68, o2293–o2294 [https://doi.org/10.1107/S1600536812028309]

4-(Prop-2-yn-1-yloxy)benzene-1,2-dicarbonitrile

Yee Jan Chin, Ai Ling Tan, Franz L. Wimmer, Aminul Huq Mirza, David J. Young, Seik Weng Ng and Edward R. T. Tiekink

S1. Comment

Phthalocyanine dyes can be made soluble in water and organic solvents by addition of suitable alkoxy or aryloxy groups (Jiang *et al.*, 2011; Sleven *et al.*, 2001). This is most easily achieved from the correspondingly substituted phthalonitriles which, in turn, are either prepared by Sandmeyer reaction of alkyl or alkoxy functionalized dihalobenzenes (Li & Lieberman, 2001; Sleven *et al.*, 2001) or aryloxy / alkoxy displacement of the corresponding halophthalonitrile (Wöhrle *et al.*, 1993; Li *et al.*, 2008; Foo *et al.*, 2012) or 4-nitrophthalonitrile (Wu *et al.*, 1998; Seven *et al.*, 2009). The latter method is most suitable for preparing 4-alkoxyphthalonitriles and was used for preparing the title compound, 4-(prop-2-ynloxy)phthalonitrile (I).

In (I), Fig. 1, the complete molecule is generated by the application of 2-fold symmetry; the molecule is disordered about this axis. The O1 and C1 atoms lie $-0.067(3)$ and $0.059(2)$ Å out of the plane through the benzene ring, respectively. The prop-2-yn-1-yl is twisted out of the plane of the benzene ring as seen in the value of the C4—O1—C5—C6 torsion angle of $173.1(3)^\circ$ and is orientated in the opposite direction to the nitrile substituents.

In the crystal packing, supramolecular chains along the *a* axis feature owing to C—H \cdots N interactions, Table 1, and 10-membered $\{\cdots\text{HC}_3\text{N}\}_2$ synthons, Fig. 2. Chains are connected into stacks along the *c* axis by π — π interactions between the benzene rings [inter-centroid distance = $3.6978(6)$ Å = length of the *c* axis]. The layers inter-digitate along the *b* axis with no specific intermolecular interactions between them.

S2. Experimental

The title compound was prepared by modification of literature procedures (Wu *et al.*, 1998; Seven *et al.*, 2009). Under a nitrogen atmosphere, anhydrous potassium carbonate (1.12 g, 8.1 mmol) was added in two portions at 1 h intervals to a solution of propargyl alcohol (1.5 ml, 26.0 mmol) and 4-nitrophthalonitrile (0.70 g, 4.04 mmol) in dry *N,N*-dimethylformamide (7 ml). After 96 h, the crude reaction mixture was poured into water (140 ml). The green precipitate was collected by vacuum filtration, washed with water and dried. The crude product was purified by silica gel column chromatography using dichloromethane as eluent to provide 0.4 g (63.9%) of a faintly coloured solid that was recrystallized from CH_2Cl_2 / hexane as colourless prisms. Melting point = 383 K. IR ν/cm^{-1} : 3287, 3119, 3077, 2231, 2135, 1596, 1494, 1321, 1260. ^1H NMR 400 MHz (CDCl_3) δ : 7.75 (1H, d), 7.35 (1H, s), 7.28 (1H, d), 4.80 (2H, s), 2.62 (1H, s).

S3. Refinement

With the exception of the acetylenic H-atom which was refined freely, carbon-bound H-atoms were placed in calculated positions [$\text{C—H} = 0.95\text{--}0.99$ Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The molecule is disordered over a 2-fold rotation axis in an exact 1:1 ratio. The anisotropic displacement

parameters of the O1 and C4 atoms were tightly restrained to be nearly isotropic.




Figure 1

The molecular structure of (I) showing displacement ellipsoids at the 50% probability level. The molecule is disordered about the 2-fold axis - only one orientation is shown.




Figure 2

A view of the supramolecular chain along the *a* axis in (I). The C—H...N and interactions are shown as blue dashed lines.




Figure 3

A view of the supramolecular layer in the *ac* plane in (I). The C—H...O and π — π interactions are shown as blue and purple dashed lines, respectively.

4-(Prop-2-yn-1-yloxy)benzene-1,2-dicarbonitrile

Crystal data

$C_{11}H_6N_2O$

$M_r = 182.18$

Monoclinic, $C2/m$

Hall symbol: $-C\ 2y$

$a = 11.4809\ (9)\ \text{\AA}$

$b = 22.2091\ (16)\ \text{\AA}$

$c = 3.6978\ (6)\ \text{\AA}$

$\beta = 91.304\ (10)^\circ$

$V = 942.62\ (18)\ \text{\AA}^3$

$Z = 4$

$F(000) = 376$

$D_x = 1.284\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 938 reflections

$\theta = 3.3\text{--}27.5^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Prism, colourless

$0.15 \times 0.05 \times 0.05\ \text{mm}$

Data collection

Agilent SuperNova Dual

diffractometer with an Atlas detector

Radiation source: fine-focus sealed tube

Mirror monochromator

Detector resolution: $10.4041\ \text{pixels mm}^{-1}$

ω scan

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.792$, $T_{\max} = 1.000$

3258 measured reflections

1114 independent reflections

840 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.038$

$\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 3.3^\circ$

$h = -14 \rightarrow 14$

$k = -26 \rightarrow 28$

$l = -4 \rightarrow 4$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.059$

$wR(F^2) = 0.163$

$S = 1.08$

1114 reflections

86 parameters

12 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0718P)^2 + 0.8054P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.41\ \text{e \AA}^{-3}$

$\Delta\rho_{\min} = -0.28\ \text{e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.1308 (2)	0.45076 (11)	0.1768 (8)	0.0273 (7)	0.50
N1	0.15188 (16)	0.19174 (9)	0.2675 (6)	0.0376 (6)	
C1	0.10982 (16)	0.23658 (9)	0.1829 (6)	0.0260 (5)	
C2	0.05529 (16)	0.29272 (8)	0.0857 (5)	0.0219 (5)	
C3	0.1103 (2)	0.34676 (10)	0.1663 (6)	0.0349 (6)	
H3	0.1853	0.3469	0.2801	0.042*	
C4	0.0555 (3)	0.40028 (10)	0.0803 (7)	0.0446 (7)	
H4	0.0936	0.4374	0.1309	0.054*	0.50
C5	0.2499 (3)	0.44609 (17)	0.3108 (13)	0.0274 (10)	0.50
H5A	0.2962	0.4211	0.1458	0.033*	0.50
H5B	0.2522	0.4272	0.5535	0.033*	0.50
C6	0.2980 (3)	0.5082 (7)	0.3303 (12)	0.030 (3)	0.50
C7	0.3440 (4)	0.5544 (2)	0.3459 (17)	0.0463 (14)	0.50
H7	0.386 (5)	0.595 (3)	0.363 (16)	0.054 (16)*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0258 (13)	0.0187 (12)	0.0370 (15)	0.0001 (10)	−0.0069 (11)	−0.0008 (11)
N1	0.0332 (10)	0.0410 (11)	0.0383 (12)	0.0126 (9)	−0.0029 (8)	0.0074 (9)
C1	0.0191 (9)	0.0345 (12)	0.0242 (11)	−0.0003 (8)	−0.0019 (8)	0.0004 (8)
C2	0.0206 (10)	0.0234 (10)	0.0217 (10)	−0.0009 (7)	−0.0001 (8)	−0.0007 (7)
C3	0.0401 (12)	0.0399 (13)	0.0251 (12)	−0.0182 (10)	0.0071 (9)	−0.0077 (9)
C4	0.0728 (15)	0.0262 (10)	0.0355 (13)	−0.0161 (10)	0.0172 (11)	−0.0063 (9)
C5	0.0220 (19)	0.0167 (19)	0.043 (3)	0.0005 (17)	−0.0104 (17)	−0.0001 (16)
C6	0.0262 (16)	0.013 (9)	0.051 (2)	−0.004 (2)	−0.0130 (15)	−0.001 (2)
C7	0.033 (3)	0.027 (2)	0.078 (4)	0.001 (2)	−0.019 (2)	−0.001 (2)

Geometric parameters (\AA , $^\circ$)

O1—C5	1.447 (5)	C4—C4 ⁱ	1.393 (6)
O1—C4	1.455 (3)	C4—H4	0.9500
N1—C1	1.147 (3)	C5—C6	1.487 (14)
C1—C2	1.437 (3)	C5—H5A	0.9900
C2—C3	1.385 (3)	C5—H5B	0.9900

C2—C2 ⁱ	1.406 (4)	C6—C7	1.155 (16)
C3—C4	1.379 (3)	C7—H7	1.02 (6)
C3—H3	0.9500		
C5—O1—C4	125.5 (3)	C3—C4—H4	119.8
N1—C1—C2	178.4 (2)	C4 ⁱ —C4—H4	119.8
C3—C2—C2 ⁱ	119.96 (13)	O1—C5—C6	107.3 (3)
C3—C2—C1	120.26 (18)	O1—C5—H5A	110.3
C2 ⁱ —C2—C1	119.77 (10)	C6—C5—H5A	110.3
C4—C3—C2	119.6 (2)	O1—C5—H5B	110.3
C4—C3—H3	120.2	C6—C5—H5B	110.3
C2—C3—H3	120.2	H5A—C5—H5B	108.5
C3—C4—C4 ⁱ	120.43 (14)	C7—C6—C5	174.6 (7)
C3—C4—O1	110.0 (2)	C6—C7—H7	178 (3)
C4 ⁱ —C4—O1	129.56 (15)		
C2 ⁱ —C2—C3—C4	0.6 (4)	C5—O1—C4—C3	5.5 (5)
C1—C2—C3—C4	-178.2 (2)	C5—O1—C4—C4 ⁱ	-173.5 (4)
C2—C3—C4—C4 ⁱ	1.2 (4)	C4—O1—C5—C6	173.1 (3)
C2—C3—C4—O1	-177.9 (2)		

Symmetry code: (i) $-x, y, -z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C3—H3 \cdots N1 ⁱⁱ	0.95	2.62	3.509 (3)	155

Symmetry code: (ii) $-x+1/2, -y+1/2, -z+1$.