

Age-dependent decline of association between obesity and coronary heart disease: a cohort study in a remote Australian Aboriginal community

Wang, Zhiqiang; Hoy, Wendy

Published in:
BMJ Open

DOI:
[10.1136/bmjopen-2013-004042](https://doi.org/10.1136/bmjopen-2013-004042)

Published: 01/01/2013

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Wang, Z., & Hoy, W. (2013). Age-dependent decline of association between obesity and coronary heart disease: a cohort study in a remote Australian Aboriginal community. *BMJ Open*, 3(11), 1-5. [e004042].
<https://doi.org/10.1136/bmjopen-2013-004042>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

BMJ Open Age-dependent decline of association between obesity and coronary heart disease: a cohort study in a remote Australian Aboriginal community

Zhiqiang Wang, Wendy E Hoy

To cite: Wang Z, Hoy WE. Age-dependent decline of association between obesity and coronary heart disease: a cohort study in a remote Australian Aboriginal community. *BMJ Open* 2013;3:e004042. doi:10.1136/bmjopen-2013-004042

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-004042>).

Received 16 September 2013
Revised 25 October 2013
Accepted 1 November 2013



CrossMark

Centre for Chronic Disease,
School of Medicine,
University of Queensland,
Herston, Queensland,
Australia

Correspondence to
Dr Zhiqiang Wang;
z.wang@uq.edu.au

ABSTRACT

Objective: To determine whether the association between obesity and coronary heart disease (CHD) in Aboriginal adults depends on age.

Design, setting and participants: A cohort study with up to 20 years of follow-up of 849 participants aged 18–76 years in a remote Aboriginal community in the Northern Territory of Australia.

Main outcome measures: Newly diagnosed CHD cases were identified through hospital records according to ICD codes during the follow-up period. Cox proportional hazard model was used to assess whether the association between obesity and CHD depends on age.

Results: During the follow-up period, 171 participants were diagnosed with CHD. On an average, the incidence rate of CHD increased with the increasing baseline BMI, 11.3%, 16.3% and 20.2% for normal weight, overweight and obese groups, respectively. HR of CHD for obesity were 2.6 (95% CI 1.1 to 6.3), 1.2 (0.7 to 2.0) and 0.5 (0.1 to 2.1) for those <40, 40–59 and 60+ years, respectively. HRs corresponding to 1 SD increase in BMI were 1.4 (1.0 to 2.0), 1.2 (1.0 to 1.5) and 0.8 (0.5 to 1.2) for those <40, 40–59 and 60+ years, respectively. The interaction terms between age and BMI as category variables or as a continuous variable were statistically significant.

Conclusions: The association between obesity and CHD is stronger for younger adults than for older adults in Aboriginal Australians in the remote community. Our findings suggest that weight control efforts may produce more beneficial effects in CHD prevention in young adults than in older adults.

INTRODUCTION

It has been well established that overweight and obesity increase the risk of mortality and coronary heart disease (CHD) in a number of studies over several decades.^{1–8} Weight reduction has been found to be beneficial in reducing the CHD risk among overweight and obese people.^{9–11} It has been suggested the effect of body mass index (BMI) on

Strengths and limitations of this study

- Major strength of the study is the long-term follow-up in a unique high-risk remote Aboriginal community.
- However, since body mass index (BMI) was estimated only once at one age point for each participant, the effect of BMI changes on CHD risk could not be directly assessed.
- Another limitation is that the generalisability of the findings needs to be further verified as this study was conducted in a single Aboriginal community.

cardiovascular disease mortality is modified by age.¹² A recent analysis by the Emerging Risk Factors Collaboration of a database of over 200 000 participants in 51 studies from 17 countries shows that the strength of the association between BMI and CHD was stronger for younger adults than for older adults.¹³ On the other hand, a study on obesity and mortality in the USA suggested that the diminishing obesity–mortality association with age was confounded by age at survey and cohort effects, and once those factors were accounted for, the obesity–mortality became stronger with age.¹⁴ Understanding the presence of such effect modification by age is important for directing prevention efforts and prioritising public health education. Aboriginal Australians have a high risk of CHD and their CHD is diagnosed at a younger age than non-Aboriginal people.¹⁵ It is still not known whether obesity has different effects on CHD risk among Aboriginal people of different ages. The objective of this study was to determine whether age modifies the association between BMI and CHD risk. We analysed the cohort data with up to 20 years follow-up from a remote Aboriginal community in the Northern Territory of Australia.

METHODS

Participants and CHD events

This is a prospective cohort study with up to 20 years of follow-up. Participants were recruited from a remote Aboriginal community in the Northern Territory of Australia from 1992 to 1995. Weight and height were measured for 849 participants aged 18–75 years and who were free from clinically apparent CHD at baseline, representing over 80% of those within the age range in the region. Those participants were followed up until 31 May 2012. During the follow-up period, new CHD events were identified through hospital records using the International Classification of Diseases codes (ICD 9 codes 410–414, and ICD 10 codes I20–I25), including myocardial infarction (410, I21), angina pectoris (411, I20) and other ischaemic heart disease (413, 414, I22, I23, I24 and I25). Deaths and their causes during the follow-up period were determined through a list of death records maintained at the community clinics. For those participants who reached a CHD event or died from non-CHD causes during the follow-up, their follow-up time was the period from the age of their initial screening visit to the age of the first CHD event or death. Others who survived the follow-up period were censored at 31 May 2012. Because individual hospital registration numbers which we used to track study participants were unique throughout the Northern Territory, we were able to capture their hospitalisation records even if our study participants had moved outside the local region. The chance of being hospitalised outside the Northern Territory was extremely low, if any, for people in this remote isolated region. Those without hospital and death records were regarded as free from CHD during the follow-up period.

Baseline BMI measurements

Height and weight were measured for study participants at baseline during 1992–1995. Height was measured to the nearest 0.5 cm without shoes. Weight was measured to the nearest 0.1 kg with participants wearing light

clothes only and without shoes. BMI was defined as weight divided by height squared (kg/m^2). Since those measurements had been taken several years before the development of CHD, BMI readings were not biased by the presence of CHD endpoints.

Statistical analysis

The data were partitioned into age bands of <40, 40–59 and 60+ years throughout the follow-up. For those whose age fell into two age bands during the follow-up period, their total follow-up time was subdivided and allocated into the corresponding age bands as described by Clayton and Hills.¹⁶ We calculated the incidence rates of CHD according to their baseline BMI values: <25, 25–29.9 and 30+ kg/m^2 . HRs of CHD for obesity and 1 SD increase in BMI were calculated with Cox proportional hazards models adjusting for baseline age and gender. We assessed effect-modification with a formal test of the interaction term between age and BMI (or obesity). To further control the effect of competing risk of non-CHD death, we used a competing-risks regression method¹⁷ to estimate the HRs. All analyses were performed with Stata SE V.12.1.¹⁸

RESULTS

Table 1 shows the characteristics of the study participants with different baseline BMI levels. Those obese participants were more likely to be women. They also had higher levels of systolic and diastolic blood pressure, and higher prevalence of known diabetes but lower prevalence of cigarette smoking and alcohol drinking at baseline.

During the follow-up period of 12 398 person-years in 849 participants, 128 died from non-CHD causes and 171 were diagnosed with CHD. Table 2 shows the detailed follow-up time and the number of cases by age and BMI groups. On an average, the incidence rates of CHD, which increased with the increasing baseline BMI, were 11.3%, 16.3% and 20.2% for normal weight, overweight and obese groups, respectively.

Table 1 Baseline characteristics of study participants by baseline BMI levels: mean (SD)

	Baseline BMI (kg/m^2)		
	Normal weight <25	Overweight 25–29.9	Obese ≥ 30
Number	524	200	125
Male, n (%)	285 (54.4)	100 (50.0)	40 (32.0)
Age, years	34.2 (13.1)	36.0 (11.1)	35.5 (9.5)
Weight, kg	57.0 (9.5)	74.8 (7.1)	91.1 (11.9)
Height, cm	166.3 (7.9)	165.9 (7.4)	164.8 (8.5)
BMI, kg/m^2	20.5 (2.6)	27.2 (1.3)	33.5 (3.2)
Waist c., cm	81.6 (9.5)	97.0 (7.2)	110.5 (9.4)
Systolic pressure, mm Hg	119.2 (18.5)	125.8 (18.0)	126.4 (18.9)
Diastolic pressure, mm Hg	73.0 (13.6)	77.7 (12.6)	79.7 (15.7)
Smoking, n (%)	422 (80.5)	137 (68.5)	75 (60.0)
Drinking, n (%)	328 (62.6)	117 (58.5)	53 (42.4)
Known diabetes, n (%)	46 (8.8)	25 (12.5)	23 (18.4)

Table 2 Incidence rate of CHD (per 1000 person-years) by age groups and baseline BMI values

Age (years)	Normal weight			Overweight			Obese		
	Person-years	Events	Rate	Person-years	Events	Rate	Person-years	Events	Rate
<40	4354	16	3.7	1249	8	6.4	742	9	12.1
40–59	2795	52	18.6	1446	31	21.4	925	25	27.0
60+	526	19	36.1	247	9	36.4	115	2	17.3
Total	7675	87	11.3	2942	48	16.3	1782	36	20.2

A rising trend in CHD risk with increased BMI was observed in younger age groups (figure 1). Estimated HRs with adjustment of baseline age, gender, known diabetes and smoking status are shown in table 3. For those younger than 40 years, adjusted HRs for overweight and obesity were 1.5 (95% CI 0.6 to 3.5) and 2.6 (95% CI 1.1 to 6.3), respectively, with BMI <25 kg/m² as the reference group. For those aged 40–59 years, their HRs of CHD were 1.1 (95% CI 0.7 to 1.8) and 1.2 (95% CI 0.7 to 2.0) for overweight and obesity, respectively. However, for those aged 60 years or older, the CHD risk for normal BMI group was as high as those for overweight and obese groups and no increasing trend in CHD risk with increasing BMI was observed in this age group. To assess whether the association between BMI and CHD risk depended on age and gender, we tested the interaction terms of BMI with age and gender in the Cox proportional hazard models. The overall test for interaction terms between age and BMI categories was statistically significant (p=0.012).

When BMI was treated as a continuous variable, the HRs corresponding to 1 SD increase in BMI decreased with increasing age, 1.4, 1.2 and 0.8 for those younger

than 40, 40–59 and 60 years or older, respectively. The interaction term between age and BMI as a continuous variable was also statistically significant (p=0.011).

Further adjusting for competing risk of non-CHD deaths, the pattern of the association between CHD and BMI either as a continuous variable or as categorical variables remained similar (table 4). The association was stronger for younger adults than for older adults with a statistically significant interaction term between age and BMI.

DISCUSSION

In this cohort study with up to 20 years follow-up, we found that age is a strong effect-modifier for the association between obesity (or BMI) and CHD in Aboriginal people. The association between obesity and CHD declined with age. The younger they are, the stronger the association between BMI and CHD. Among adults under 40 years, the risk of CHD is 2.6 times in obese individuals as in those with BMI <25 kg/m², but there is no significant association between obesity and CHD among those 60 years or older. Our findings suggest that the effect of obesity on development of CHD is stronger in younger people than in older people.

It has been well established that the people with high BMI have a high risk of CHD,^{2–8} but a very few studies have assessed the effect modification of the BMI-CHD association by age. The Emerging Risk Factors Collaboration group suggested that age is a strong effect modifier for BMI-CHD association.¹³ They reported that the HRs for CHD associated with per 1 SD increase in BMI decreased with increasing age, from 1.41 for their youngest group (40–59 years) to 1.23 for 60–69 and 1.12 for 70 years or older.¹³ In our study, we used different age groups because CHD occurred at younger ages in Aboriginal people.¹⁵ Nevertheless, the pattern of effect modification by age in our study was similar to that reported by the Emerging Risk Factors Collaboration group. To our knowledge, this is the first study to assess effect modification by age in a high-risk indigenous population.

It has also been reported that excess body weight increases the risk of death from any cause and from cardiovascular disease and the relative risk with greater body weight is higher among younger participants in a study of US adults.^{12 19} However, a study by Masters *et al* suggested that survey-based estimates of age patterns in



Figure 1 Incidence of coronary heart disease by age and body mass index in Aboriginal adults.

Table 3 HRs* of CHD for different baseline BMI levels by age groups

Age (years)	Overweight† HR (95% CI)	Obese† HR (95% CI)	Interaction p Value‡	BMI (1 SD) HR (95% CI)	Interaction p Value§
<40	1.5 (0.6 to 3.5)	2.6 (1.1 to 6.3)	0.012	1.4 (1.0 to 2.0)	0.011
40–59	1.1 (0.7 to 1.8)	1.2 (0.7 to 2.0)		1.2 (1.0 to 1.5)	
60+	1.0 (0.5 to 2.5)	0.5 (0.1 to 2.1)		0.8 (0.5 to 1.2)	

*Adjusted for age, gender, known diabetes and smoking status.

†With BMI <25 kg/m² as reference.

‡Interaction between age and BMI as category variables.

§Interaction between age and BMI as a continuous variable.

obesity–mortality relationship are confounded by age at survey and cohort membership. When those factors are accounted for, Masters *et al*¹⁴ concluded that the obesity–mortality relationship becomes stronger with age. However, their conclusions might be due to the misinterpretation of main effect coefficients in the presence of interactions.²⁰ In this study, controlling for baseline age did not alter in obesity–mortality relationship in different age groups. Those biases caused by cohort membership that Masters *et al* proposed were unlikely present because this study was based on the data from a single cohort. Adjusting for the competing risk of non-CHD death and the potential confounding effects of smoking, diabetes and gender did not alter the age pattern of the BMI–CHD relationship.

It is not clear why the effect of obesity (or higher BMI) on development of CHD declines as people get older. A similar pattern of the age-dependent significant attenuation between obesity and CHD risk factors such as hyperglycaemia, hypertension and dyslipidaemia could partly explain the observed age-dependent decline between obesity and development of CHD.^{21–22} This phenomenon, if further confirmed in interventional studies, may have clinical and public health implications of weight control in people of different ages. Our findings stress the importance of weight reduction targeting younger adults for reducing CHD risk in this population. The same amount of weight reduction in younger people may produce more beneficial effects on CHD risk reduction than in their older counterparts. Several other factors have been found to be associated with CHD or cardiovascular disease risk in this population such as C reactive protein²³ and albuminuria²⁴ in addition to conventional risk factors.^{25–26} Since obesity

has been considered to affect CHD risk through its influence on those known risk factors,²⁷ we did not adjust for those known risk factors in our analysis.

There are several limitations in this study. BMI was estimated only once at one age point for each participant and no recent BMI values were taken. Therefore, the effect of BMI changes on CHD risk could not be directly assessed. The CHD cases were identified through routinely documented hospital records during the follow-up period. Under-reporting is possible as some participants with minor CHD events may not be hospitalised and diagnosed as such. However, there is no evidence that such under-reporting could have occurred differently among people with different BMI levels. Another limitation of this study is the small sample size, particularly for the older age group. Although the interaction term for testing the dependence of the BMI–CHD association on age was statistically significant, the association between BMI and CHD in the older age group could not be accurately estimated as reflected by the wide 95% CIs. Finally, the study was conducted in a single relatively homogenous population in one community. It remains to be verified whether our findings can be applied to the general Aboriginal population in Australia.

CONCLUSION

The association between obesity and CHD declines with increasing age and is stronger in young adults than for older adults in Aboriginal Australians. Our findings imply that the efforts of obesity prevention and weight control are more likely to produce beneficial effects on CHD prevention in younger than in older people. Further interventional studies are needed to quantify

Table 4 HRs of CHD for different baseline BMI levels adjusting for competing risk of non-CHD death

Age (years)	Overweight* HR (95% CI)	Obese* HR (95% CI)	Interaction p Value†	BMI (1 SD)* HR (95% CI)	Interaction p Value‡
<40	1.5 (0.6 to 3.6)	2.6 (1.1 to 6.0)	0.026	1.4 (1.1 to 2.0)	0.012
40–59	1.2 (0.7 to 1.8)	1.3 (0.8 to 2.1)		1.2 (1.0 to 1.4)	
60+	1.2 (0.6 to 2.5)	0.5 (0.1 to 2.1)		0.8 (0.5 to 1.2)	

*Adjusted for competing risk of non-CHD death, age, gender, known diabetes and smoking status.

†Interaction between age and BMI as category variables.

‡Interaction between age and BMI as a continuous variable.

the effect of weight reduction on CHD risk in people of different ages.

Acknowledgements The authors especially thank the Aboriginal people who participated in this study. Shuqin Li at the Northern Territory Department of Health assisted in the interpretation of hospital data.

Contributors WEH and ZW conceived the idea of the study and were responsible for the design of the study. WEH provided input into the data analysis and was responsible for the acquisition of the baseline data. ZW was responsible for linking baseline and hospital data and for undertaking data analysis. Both WEH and ZW contributed to the first draft, and read and approved the final version.

Funding This work was supported by NHMRC of Australia (APP1025300/1025350).

Competing interests WEH was supported by Australian Fellowship (#511081) and ZW by NHMRC Fellowship (APP1042343).

Ethics approval The project was approved by the University of Queensland Behavioural and Social Science Ethical Review Committee (#2011001232).

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

1. Flegal KM, Kit BK, Orpana H, *et al.* Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA* 2013;309:71–82.
2. Harris TB, Launer LJ, Madans J, *et al.* Cohort study of effect of being overweight and change in weight on risk of coronary heart disease in old age. *BMJ* 1997;314:1791–4.
3. Manson JE, Colditz GA, Stampfer MJ, *et al.* A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med* 1990;322:882–9.
4. Keys A, Aravanis C, Blackburn H, *et al.* Coronary heart disease: overweight and obesity as risk factors. *Ann Intern Med* 1972;77:15–27.
5. Logue J, Murray HM, Welsh P, *et al.* Obesity is associated with fatal coronary heart disease independently of traditional risk factors and deprivation. *Heart* 2011;97:564–8.
6. Nguyen NT, Nguyen XM, Wooldridge JB, *et al.* Association of obesity with risk of coronary heart disease: findings from the National Health and Nutrition Examination Survey, 1999–2006. *Surg Obes Relat Dis* 2010;6:465–9.
7. Nanchahal K, Morris JN, Sullivan LM, *et al.* Coronary heart disease risk in men and the epidemic of overweight and obesity. *Int J Obes (Lond)* 2005;29:317–23.
8. Stevens J, Erber E, Truesdale KP, *et al.* Long- and short-term weight change and incident coronary heart disease and ischemic stroke: the Atherosclerosis Risk in Communities Study. *Am J Epidemiol* 2013;178:239–248.
9. Sartorio A, Lafortuna CL, Marinone PG, *et al.* Short-term effects of two integrated, non-pharmacological body weight reduction programs on coronary heart disease risk factors in young obese patients. *Diabetes Nutr Metab* 2003;16:262–5.
10. Vogel JA, Franklin BA, Zalesin KC, *et al.* Reduction in predicted coronary heart disease risk after substantial weight reduction after bariatric surgery. *Am J Cardiol* 2007;99:222–6.
11. Eilat-Adar S, Eldar M, Goldbourt U. Association of intentional changes in body weight with coronary heart disease event rates in overweight subjects who have an additional coronary risk factor. *Am J Epidemiol* 2005;161:352–8.
12. Stevens J, Cai J, Pamuk ER, *et al.* The effect of age on the association between body-mass index and mortality. *N Engl J Med* 1998;338:1–7.
13. The Emerging Risk Factors Collaboration. Separate and combined associations of body-mass index and abdominal adiposity with cardiovascular disease: collaborative analysis of 58 prospective studies. *Lancet* 2011;377:1085–95.
14. Masters RK, Powers DA, Link BG. Obesity and US mortality risk over the adult life course. *Am J Epidemiol* 2013;177:431–42.
15. Wang Z, Hoy WE. Lifetime risk of developing coronary heart disease in Aboriginal Australians: a cohort study. *BMJ Open* 2013;3:e002308.
16. Clayton D, Hills M. *Statistical models in epidemiology*. Oxford: Oxford University Press, 1993.
17. Fine JP, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc* 1999;94:496–509.
18. Stata Statistical Software: Release 12 [program]. College Station, TX: StataCorp LP, 2011.
19. Singh PN, Lindsted KD. Body mass and 26-year risk of mortality from specific diseases among women who never smoked. *Epidemiology* 1998;9:246–54.
20. Wang Z. Obesity and US mortality risk over adult life course: misinterpretation of main effect estimates in the presence of interactions. *Am J Epidemiol* 2013;in press.
21. Wakabayashi I. Age-dependent influence of gender on the association between obesity and a cluster of cardiometabolic risk factors. *Gen Med* 2012;9:267–77.
22. Wakabayashi I, Daimon T. Age-dependent decline of association between obesity and hyperglycemia in men and women. *Diabetes Care* 2012;35:175–7.
23. Wang Z, Hoy WE. C-reactive protein: an independent predictor of cardiovascular disease in Aboriginal Australians. *Aust N Z J Public Health* 2010;34(Suppl 1):S25–9.
24. Wang Z, Hoy WE. Albuminuria and incident coronary heart disease in Australian Aboriginal people. *Kidney Int* 2005;68:1289–93.
25. Wang Z, Hoy WE. Association between diabetes and coronary heart disease in Aboriginal people: are women disadvantaged? *Med J Aust* 2004;180:508–11.
26. Wang Z, Hoy WE. Is the Framingham coronary heart disease absolute risk function applicable to Aboriginal people? *Med J Aust* 2005;182:66–9.
27. Poirier P, Giles TD, Bray GA, *et al.* Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss: an update of the 1997 American Heart Association Scientific Statement on Obesity and Heart Disease from the Obesity Committee of the Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 2006;113:898–918.