

Correction to

Far eastern curlew and whimbrel prefer flying low - wind support and good visibility appear only secondary factors in determining migratory flight altitude (Movement Ecology, (2021), 9, 1, (32), 10.1186/s40462-021-00267-5)

Galtbalt, Batbayar; Lilleyman, Amanda; Coleman, Jonathan T.; Cheng, Chuyu; Ma, Zhijun; Rogers, Danny I.; Woodworth, Bradley K.; Fuller, Richard A.; Garnett, Stephen T.; Klaassen, Marcel

Published in:
Movement Ecology

DOI:
[10.1186/s40462-021-00289-z](https://doi.org/10.1186/s40462-021-00289-z)

Published: 01/12/2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Galtbalt, B., Lilleyman, A., Coleman, J. T., Cheng, C., Ma, Z., Rogers, D. I., Woodworth, B. K., Fuller, R. A., Garnett, S. T., & Klaassen, M. (2021). Correction to: Far eastern curlew and whimbrel prefer flying low - wind support and good visibility appear only secondary factors in determining migratory flight altitude (Movement Ecology, (2021), 9, 1, (32), 10.1186/s40462-021-00267-5). *Movement Ecology*, 9(1), Article 51. <https://doi.org/10.1186/s40462-021-00289-z>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CORRECTION

Open Access



Correction to: Far eastern curlew and whimbrel prefer flying low - wind support and good visibility appear only secondary factors in determining migratory flight altitude

Batbayer Galtbalt^{1*} , Amanda Lilleyman², Jonathan T. Coleman³, Chuyu Cheng⁴, Zhijun Ma⁴, Danny I. Rogers^{5,6}, Bradley K. Woodworth⁷, Richard A. Fuller⁷, Stephen T. Garnett² and Marcel Klaassen^{1,8}

Correction to: *Movement Ecology* (2021) 9:32 <https://doi.org/10.1186/s40462-021-00267-5>

Following publication of the original article [1], errors were identified in the presentation of some of the reference citations in the Background section (page 2 of the PDF) and the Conclusion section (page 10 of the PDF) due to a typesetting mistake.

These errors have been corrected and the original article has been updated. The publisher apologises to the authors and readers for the inconvenience caused by this mistake.

Author details

¹Centre for Integrative Ecology, School of Life and Environmental Science, Deakin University, Geelong, Victoria, Australia. ²Threatened Species Recovery Hub, National Environment Science Program, Research Institute for Environment and Livelihoods, Charles Darwin University, Ellengowan Drive, Casuarina, Northern Territory 0909, Australia. ³Queensland Wader Study Group, 22 Parker Street, Shailer Park 4128, Australia. ⁴Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Coastal Ecosystems Research Station of the Yangtze River Estuary, Institute of Biodiversity Science, School of Life Sciences, Fudan University, Shanghai 200433, China. ⁵Department of Environment, Water, Land and Planning, Arthur Rylah Institute, PO Box 137, Heidelberg, Victoria 3084, Australia. ⁶Australasian Wader Studies

Group, Melbourne, Victoria, Australia. ⁷School of Biological Sciences, University of Queensland, Brisbane, Queensland, Australia. ⁸Victorian Wader Study Group, Melbourne, Victoria, Australia.

Published online: 20 October 2021

Reference

1. Galtbalt, et al. Far eastern curlew and whimbrel prefer flying low - wind support and good visibility appear only secondary factors in determining migratory flight altitude. *Mov Ecol.* 2021;9:32. <https://doi.org/10.1186/s40462-021-00267-5>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1186/s40462-021-00267-5>.

*Correspondence: batbr19@gmail.com

¹ Centre for Integrative Ecology, School of Life and Environmental Science, Deakin University, Geelong, Victoria, Australia
Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.